



UNIFORMED SERVICES UNIVERSITY OF THE HEALTH SCIENCES
F. EDWARD HÉBERT SCHOOL OF MEDICINE
4301 JONES BRIDGE ROAD
BETHESDA, MARYLAND 20814-4799



GRADUATE PROGRAMS IN THE
BIOMEDICAL SCIENCES AND
PUBLIC HEALTH

May 21, 2009

APPROVAL SHEET

Ph.D. Degrees

Interdisciplinary

- Emerging Infectious Diseases
- Molecular & Cell Biology
- Neuroscience

Departmental

- Clinical Psychology
- Environmental Health Sciences
- Medical Psychology
- Medical Zoology
- Pathology

Doctor of Public Health (Dr.P.H.)

Physician Scientist (MD/Ph.D.)

Master of Science Degrees

- Public Health

Masters Degrees

- Military Medical History
- Public Health
- Tropical Medicine & Hygiene

Graduate Education Office

Dr. Eleanor S. Metcalf, Associate Dean
Bettina Arnett, Support Specialist
Roni Bull, Support Specialist

Web Site

<http://www.usuhs.mil/graded/>

E-mail Address

graduateprogram@usuhs.mil


Phone Numbers

Commercial: 301-295-9474
Toll Free: 800-772-1747
DSN: 295-9474
FAX: 301-295-6772


Title of Dissertation: "Branched-chain Amino and Keto Acid Biochemistry and Cellular Biology in Central Nervous System Diseases"

Name of Candidate: Jeremy Henriques
Doctor of Philosophy Degree
30 March 2009


Dissertation and Abstract Approved:


Merrily Poeth, M.D.
Department of Pediatrics
Committee Chairperson


5/22/09
Date


Ajay Verma, M.D., Ph.D.
Department of Neurology
Committee Member


Date


Michael Schell, Ph.D.
Department of Pharmacology
Committee Member


May 22, 2009
Date


Saibal Dey, Ph.D.
Department of Biochemistry
Committee Member


May 22, 2009
Date


Timothy O'Neill, Ph.D.
Department of Pediatrics
Committee Member

May 22, 2009
Date


Jay Thakar, Ph.D.
Department of Neurology
Committee Member

May 22, 2009
Date


Gregory P. Mueller
Department of Anatomy, Physiology
and Genetics
Committee Member

May 22, 2009

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 21 MAY 2009		2. REPORT TYPE		3. DATES COVERED 00-00-2009 to 00-00-2009	
4. TITLE AND SUBTITLE Branched-Chain Amino And Keto Acid Biochemistry And Cellular Biology In An In Vitro Model Of Central Nervous System Diseases				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Uniformed Services University of The Health Services,4301 Jones Bridge Rd,Bethesda,MD,20814				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Branched-chain amino acids (BCAAs: leucine, isoleucine, and valine) are essential components of many biochemical and biological processes. There are well-established pathways, such as fatty acid synthesis and oxidation, anabolic use to synthesize new proteins, and anaplerotic use to generate or sustain metabolic intermediate molecules, which we define as "classical" pathways here. Recently new "non-classical" roles for BCAAs have been discovered, most notably for leucine. Leucine has been shown to initiate protein synthesis by increasing translational protein complex activities and to suppress feeding behaviors in rats; both phenomena are at least partially mediated through the mammalian target of rapamycin (mTOR) kinase cascade. Our lab previously identified another potential non-classical pathway independent of mTOR, acting through the hypoxia inducible factor (HIF) transcriptional regulatory protein. Our goal for the first line of research for this work was to validate and further elaborate that leucine promotes HIF transcriptional activation through its 2-oxoacid derivative, 2-ketoisocaproic acid using an in vitro glioma cell line transgenically altered to express a reporter protein for HIF-1; degradative cycle activity. The 2-oxoacid was found to increase the half-life of the HIF-1; component of the HIF-1 heterodimeric complex in normal oxygen conditions and induce secretion of a known HIF-1 transcriptional protein target independent of mTOR activation.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 116	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

The author hereby certifies that the use of any copyrighted material in the thesis manuscript entitled:

"Branched-chain Amino- and Keto acid Biochemistry and Cellular Biology in an *in vitro* Model of Central Nervous System Disease"

is appropriately acknowledged and, beyond brief excerpts, is with the permission of the copyright owner.

A handwritten signature in black ink, appearing to read 'Jeremy Henriques', with a long, sweeping horizontal line extending to the right.

Jeremy Henriques
Neuroscience Program
Uniformed Services University

**Branched-chain Amino- and Keto acid Biochemistry and Cellular Biology in an
in vitro Model of Central Nervous System Disease**

Jeremy Henriques, Doctor of Philosophy in Neuroscience, 2008

Supervised by Ajay Verma, MD, PhD, Former Associate Professor of Neuroscience

Abstract

Branched-chain amino acids (BCAAs: leucine, isoleucine, and valine) are essential components of many biochemical and biological processes. There are well-established pathways, such as fatty acid synthesis and oxidation, anabolic use to synthesize new proteins, and anaplerotic use to generate or sustain metabolic intermediate molecules, which we define as “classical” pathways here. Recently, new “non-classical” roles for BCAAs have been discovered, most notably for leucine. Leucine has been shown to initiate protein synthesis by increasing translational protein complex activities and to suppress feeding behaviors in rats; both phenomena are at least partially mediated through the mammalian target of rapamycin (mTOR) kinase cascade. Our lab previously identified another potential non-classical pathway independent of mTOR, acting through the hypoxia inducible factor (HIF) transcriptional regulatory protein. Our goal for the first line of research for this work was to validate and further elaborate that leucine promotes HIF transcriptional activation through its 2-oxoacid derivative, α -ketoisocaproic acid, using an *in vitro* glioma cell line transgenically altered to express a reporter protein for HIF- α degradative cycle activity. The 2-oxoacid was found to increase the half-life of the HIF-1 α component of the HIF-1 heterodimeric complex in normal oxygen conditions and induce secretion of a known HIF-1 transcriptional protein target

independent of mTOR activation. The effect of 2-oxoacid stimulation of HIF-1 transcriptional activation was inhibited by addition of ascorbic acid, a cofactor known to increase HIF degradation in normoxic conditions. Our goal for the second line of research was to investigate the interaction between classical and non-classical BCAA pathways, that is to say the interaction between metabolic and signaling roles in an *in vitro* model of cancer. BCAAs exclusively undergo irreversible decarboxylation and oxidation through the branched-chain α -ketoacid dehydrogenase complex (BCKDC), a multienzyme complex regulated by reversible phosphorylation. By knocking down the kinase, which normally decreases BCKDC activity, with a stable shRNA transfection in a glioblastoma cell line, the BCKDC exhibited decreased activity and phosphorylation signal compared to wild-type and control vector cell lines. We expected knockdown of the kinase would decrease the anabolic protein translational pathway to result in a decrease of cancerous phenotypes, such as proliferation, invasion, migration, and colony formation, but found the opposite. Further investigation indicated compensatory changes to glycolytic pathways such as decreased pyruvate dehydrogenase complex (PDC) activity and increased lactate production. These characteristics are known to increase the cancerous phenotypes investigated, a phenomenon called “The Warburg Effect”. These results implicate BCAAs as an underappreciated component in cancer biochemistry, but also relate to various other lines of research, such as clinical sequelae of a congenital genetic disorder known as maple syrup urine disease (MSUD), neurotransmitter homeostasis and disruption in epilepsy, and muscle wasting in patients with cancer known as cancer cachexia.

**Branched-chain Amino- and Keto acid Biochemistry and Cellular Biology in an
in vitro Model of Central Nervous System Disease**

By

Jeremy Henriques

Dissertation submitted to the Faculty of the Program in Neuroscience of the
Uniformed Services University of the Health Sciences in partial fulfillment of the
requirements for the Degree of Doctor of Philosophy, 2008.

Acknowledgements

I would like to acknowledge the researchers who built the foundation of knowledge from which this work stems. In particular, Drs. Susan Hutson, Christopher Lynch, Nam Ho Jeoung and Robert Harris, all of whom personally collaborated with me on these projects adding invaluable services and insight. To the members of my doctoral advisory committee, I truly appreciate and am forever indebted to your collective assistance. To the USUHS faculty and staff, thank you so very much for all of your support. I have always been a challenge and cannot stress how much I appreciate all of the hard work and mentoring I obtained. In particular, I would like to thank Dr. Regina Armstrong for her guidance and patience, Dr. Gudrun Ihrke for her intellectual and technical expertise, Dr. Neil Grunberg for his scholarly mentorship, Dr. Michael Schell for his understanding, patience, and guidance through extraordinary circumstances, Dr. Merrily Poth for her undying moral and intellectual support and Dr. Ajay Verma for his mentorship in developing a scholarly career. To my friends who were (and continue to be) invaluable resources of comfort, stimulation, and support, most importantly Drs. Ahmed Mohyeldin and Thomas McFate, cohorts in the Verma Lab who will always be dear to me. Good luck to you both on your future endeavors. To my family old and new, without whom I would never have had the drive to attend college, much less complete my doctoral degree. Last and most importantly, to my wife who I met in November of my first year and has been the most important source of drive, pride, and love for me throughout my doctoral work. My life is entwined with yours, for better and worse, and I would not have it any other way.

Table of Contents

94		
95		
96	Abstract.....	ii
97	Acknowledgements	v
98	Table of Contents.....	vi
99	List of Figures	viii
100	Abbreviations	ix
101	Introduction	1
102	<i>Background.....</i>	<i>1</i>
103	<i>Rationale and Hypotheses.....</i>	<i>2</i>
104	<i>Survival: The Essence of Evolution.....</i>	<i>3</i>
105	<i>HIF biology.....</i>	<i>5</i>
106	<i>Transcriptional activation of biochemical molecules (ie, glycolytic enzymes, metabolic</i>	
107	<i>regulators).....</i>	<i>6</i>
108	<i>HIF and the Warburg Effect</i>	<i>8</i>
109	<i>Cancer hypermetabolism</i>	<i>8</i>
110	<i>Branched-chain amino acids in cancer</i>	<i>10</i>
111	<i>Branched-chain Amino Acids.....</i>	<i>11</i>
112	<i>Metabolism through the BCKDC.....</i>	<i>12</i>
113	<i>New signaling roles have emerged (the comprehensive model)</i>	<i>14</i>
114	<i>Summary and Segue</i>	<i>16</i>
115	Branched-chain α-ketoacids increase HIF-1 signaling independently of mTOR	17
116	<i>Summary.....</i>	<i>18</i>
117	<i>Introduction</i>	<i>19</i>
118	<i>Experimental Methods</i>	<i>23</i>
119	<i>Results</i>	<i>26</i>

120	<i>Discussion</i>	31
121	<i>References</i>	33
122	<i>Legends to Figures</i>	43
123	<i>Figures</i>	46
124	Alterations to branched-chain amino acid metabolism increase <i>in vitro</i> malignant	
125	characteristics of C6 glioma cells	49
126	<i>Summary</i>	50
127	<i>Experimental Methods</i>	55
128	<i>Results</i>	60
129	<i>Discussion</i>	65
130	<i>References</i>	68
131	<i>Legends to Figures</i>	73
132	<i>Figures</i>	76
133	General Discussion	82
134	References	93
135		

136 **List of Figures**

137	Figure 1. BCAAs and BCKAs stabilize HIF-1 α in normoxia, reversible by ascorbate	
138	treatment.....	46
139	Figure 2. Dose-response for BCAAs and BCKAs on HIF-1 α level stabilization and	
140	functional roles in gene upregulation	47
141	Figure 3. KIC induces HIF-1 α stabilization in primary astrocyte-enriched rat cultures,	
142	reversible by ascorbate treatment.....	48
143	Figure 4. Viability and functionality of BCKDC-kinase knockdown in vitro system...	76
144	Figure 5. Analysis of BCKDC additional subunits	77
145	Figure 6. Cell proliferation in serum and serum-free media show differences between	
146	shK cells and wild type and CVC6 control cells	78
147	Figure 7. Metastatic phenotype analyses indicate differences between shK cells and	
148	wild type and CVC6 control cells.....	79
149	Figure 8. Differential phosphorylation status of the mitochondrial metabolic complex	
150	PDC for shK and wild type and CVC6 controls	80
151		

152

Abbreviations

153 ANOVA –analysis of variance

154 ATP – adenosine 5'-triphosphate

155 BCA – bicinchoninic acid

156 BCAA – branched-chain amino acid

157 BCAT – branched-chain amino acid transaminase

158 BCKA – branched-chain keto acid

159 BCKDC – branched-chain α -keto acid dehydrogenase complex

160 BDK – branched-chain α -keto acid dehydrogenase complex kinase

161 BDP – branched-chain α -keto acid dehydrogenase complex phosphatase

162 CCD – charge-coupled device

163 CIC – α -chloroisocaproic acid

164 CMA – cylindrical mirror analyzer

165 CNS – central nervous system

166 CSF – cerebrospinal fluid

167 CVC6 – shRNA control vector C6 glioma cells

168 DMEM – Dulbecco's Modified Essential Medium

169 DNA – deoxyribonucleic acid

170 ECM – extracellular matrix

171 ELISA – enzyme-linked immunosorbent assay

172 FBS – fetal bovine serum

173 FIH – factor inhibiting HIF

174 GBM – glioblastoma multiforme

175 GFP – green fluorescent protein
176 HIF – hypoxia inducible factor
177 HPH – HIF prolyl hydroxylase
178 HSD – honest significant difference
179 KIC – α -ketoisocaproic acid
180 LAT – L-type amino acid transporter
181 MSUD- maple-syrup urine disease
182 mTOR – mammalian target of rapamycin
183 ODD – oxygen-dependent degradation domain
184 PBS- phosphate buffered saline
185 PDC – pyruvate dehydrogenase complex
186 PDK – pyruvate dehydrogenase complex kinase
187 PHD – prolyl-hydroxylase domain
188 PLSD – protected least squared difference
189 RNA – ribonucleic acid
190 S6R – S6 ribosomal protein
191 SDS – sodium dodecyl sulfate
192 shK – BDK shRNA C6 knockdown cells
193 SPSS – statistical package for the social sciences
194 TCA – tricarboxylic acid
195 tRNA – transfer RNA
196 VEGF – vascular endothelial growth factor
197

Introduction

Background

Branched-chain amino acids (BCAAs: leucine, isoleucine, and valine) are essential components of many biochemical and biological processes. Unlike many amino acids, humans are unable to synthesize BCAAs *de novo* and must obtain them through dietary consumption. This fact is important when one realizes the many different roles the BCAAs play in human physiology; some well established pathways involving BCAAs are fatty acid synthesis and oxidation, anabolic use to synthesize new proteins, and anaplerotic use to generate or sustain metabolic intermediate molecules. These functions for BCAAs are the oldest and most widely known, which we define as “classical” pathways here. Recently, new “non-classical” roles for BCAAs have been discovered, most notably for leucine. As stated above, BCAAs are utilized, like other amino acids, as building blocks for protein synthesis. When there are insufficient amounts of BCAAs available for protein synthesis, functional proteins are unable to be synthesized. Until recently it was unclear how BCAA limitation effectively terminated protein synthesis. It was discovered that leucine acts to initiate the congregation of proteins necessary for protein synthesis, collectively referred to as ‘protein translational machinery’ in this work, by interaction with multiple signal transduction and regulatory molecules (Kimball and Jefferson 2006). Along the same line of research, it was discovered that leucine interaction with signal transduction molecules occurs in the hypothalamus of rats to increase protein translation locally, and also to decrease feeding behaviors (Cota and others 2006). Both actions were inhibited by blocking one of the signaling molecules,

mammalian target of rapamycin (mTOR), downstream of leucine. It becomes evident from this prior work that leucine and the other BCAAs are important for *both* biochemical and molecular biological functions. We became interested in both aspects of BCAA function (ie, as metabolic and signaling, or signal initiation, molecules).

Rationale and Hypotheses

Normal cells grow, perform necessary and regulated functions based on expression of an appropriate set of genes, then die. Senescence and apoptosis are mechanisms through which the body regulates its own function. However, when genetic mutation leads to dysregulation of the cell cycle, the cell can lose its ability to self-regulate its growth cycle. These new, abnormal cells outlive normal cells and begin to populate tissue or an organ at a higher rate than normals cells, which can cause cancer. Half of all men and one third of all women in the US will be diagnosed with cancer within their lifetimes. Cancers can begin in many different parts of the body, but different types of cancer have varying biochemical and cytological characteristics. For example, lung cancer and breast cancer are very different diseases. They grow at different rates and respond to different treatments. That's why people with cancer need treatment that has shown to be effective for their particular type of cancer. While there have been great advances in targeted chemotherapy, investigations into basic biochemistry have led to exciting results that could have implications for many, if not all, types of cancers. If a novel molecular target can be identified that can change the way cells behave on a fundamental,

biochemical level, it could bring new insight into the therapeutic strategy against cancers.

Recently, our lab had identified pyruvate dehydrogenase complex (PDC) as one such metabolic enzyme whose activity could predictably and consistently be correlated to cancer phenotypes. In this work, we propose the branched-chain α -ketoacid dehydrogenase complex (BCKDC) as another metabolic enzyme target whose substrates and products act as signaling molecules and whose activity can be altered to change cancer phenotypes. Our hypotheses were:

1. α -ketoisocaproic acid, a substrate of the BCKDC and deaminated product of leucine, can act to decrease HIF degradation cycle activity thereby promoting neoplastic activity in an *in vitro* model of CNS cancer
2. cells genetically modified by small hairpin RNA for the BCKDC kinase will increase BCKDC activity and indicate a reciprocal relationship between BCKDC activity and cellular aggressiveness (eg, proliferation, migration, invasion, colony formation) in an *in vitro* model of CNS cancer.

Survival: The Essence of Evolution

Adaptability is the essence of evolution. Throughout the development of our modern cell, the environment has exerted pressure which allowed the emergence of advantageous traits. These changes result in progeny that are more adaptable to environmental stress. There have been hundreds of survival mechanisms discovered in the mammalian cell. Survival mechanisms also entail senescence and apoptosis, since an aberrant cell can lead to widespread dysregulation and system failure or death. When these mechanisms function normally, a cell is able to

withstand a certain threshold of stress greater than the cell would normally allow. However, when one of these mechanisms is dysregulated, the effects can be severe. Perhaps the most common life-threatening disease caused by mutations to genes encoding for regulatory proteins is cancer. Among the causes of cancer are: mutations to the so-called “tumor suppressor” genes; to the genes encoding for apoptotic pathway proteins; and to the genes encoding for cell cycle regulatory proteins. But cancer as an *in vitro* cellular phenotype and as a clinical disease is distinctly different. Much like a square is a rectangle but a rectangle is not a square, *in vitro* cancer is uncontrolled proliferation but this uncontrolled proliferation does not necessarily constitute a clinical diagnosis of cancer. Cancer disease is defined by Hanahan and Weinberg (2000) as having six hallmarks: self-sufficiency in growth signals, insensitivity to growth-inhibitory (antigrowth) signals, evasion of programmed cell death (apoptosis), limitless replicative potential, sustained angiogenesis, and tissue invasion and metastasis (Hanahan and Weinberg 2000). Recent advances have determined that there are a handful of molecules involved in many, if not all, of these phenomena seen in cancer. One such protein is hypoxia inducible factor (HIF), an evolutionarily conserved heterodimeric transcription factor.

Another commonly observed characteristic of cancer cells, that some consider the seventh hallmark of cancer, is altered glucose metabolism, which is characterized by high levels of lactate in normal oxygen conditions when compared to normal tissue. HIF is known to regulate over 100 genes encoding for growth factors, metabolic enzymes, and signaling molecules, though its evolutionary roles

has been to respond to hypoxic, or low oxygen environments. The survival response induces, in particular, a rapid switch to glycolysis and subsequent angiogenesis.

HIF biology

The hypoxic response involves an evolutionarily conserved protein known to be regulated primarily by oxygen concentrations. HIF is a protein involved in signaling survival mechanisms in cells under hypoxic, or low oxygen conditions (Goldberg and others 1988; Semenza and others 1991). HIF is a heterodimeric transcriptional regulator that upregulates the expression of over 100 genes, including glycolytic enzymes and vascular growth factors (for review (Ke and Costa 2006)). Recent estimates using DNA microarray analyses indicate that as much as 2% of human genes in arterial endothelia are regulated by HIF-1 (Manalo and others 2005). HIF-1 was discovered by the identification of a hypoxia response element (HRE) present in the gene encoding for erythropoietin, the primary factor involved in the proliferation of red blood cells (Goldberg and others 1988; Semenza and others 1991). The active HIF complex is a heterodimer comprised of an α and a β subunit. The α -subunit of HIF is constitutively degraded under normoxia, or normal oxygen conditions, by the HIF prolyl hydroxylase (HPH, or alternatively prolyl hydroxylase domain proteins or PHDs) enzymatic cycle. The HPH enzymes affix an –OH group to proline-402 or -564 resulting in ubiquitin-mediated degradation of the subunit (Masson and others 2001; Srinivas and others 1999). In hypoxic conditions, the availability of oxygen becomes a limiting factor that reduces the constitutive hydroxylation of HIF- α necessary for targeted degradation. As a result, HIF- α levels are stabilized which enables association with HIF- β , a generic transcription cohort

(Wang and others 1995). Once HIF- α/β have dimerized, the HIF complex translocates to the nucleus to associate with other transcription factors for targeted transcriptional upregulation. Another regulatory protein, factor inhibiting HIF (FIH), hydroxylates an aspartate residue in the HIF- α subunit in the nucleus in normoxic conditions. This hydroxylation prevents association with p300, an essential interaction for transcriptional activation, which inhibits hypoxia-induced gene expression (Dalgard and others 2004; Lando and others 2002a; Lando and others 2002b).

The HRE sequence previously discussed provided for the discovery of HIF (Goldberg and others 1988; Semenza and others 1991). Genes with an imbedded HRE sequence (5'-RCGTG-3')¹ are targeted by HIF upon activation, primarily by hypoxia (although other factors have been shown to activate HIF [for review (Ke and Costa 2006)]). One such encoding sequence is the VEGF gene (Levy and others 1995).

Transcriptional activation of biochemical molecules (ie, glycolytic enzymes, metabolic regulators)

As stated in the previous section, HIF activation has been shown to upregulate the expression of over 100 transcripts (for review (Ke and Costa 2006)). Not only are molecular alterations necessary for protection and survival, but metabolism must also be affected in order to sustain physiological function. HIF is a known regulator of enzymes involved in glycolysis. Glucose transporter-1 and -3

¹ R = G or A

332 (Chen and others 2001), lactate dehydrogenase-A (Semenza and others 1996),
333 glyceraldehyde phosphate dehydrogenase (Graven and others 1999), hexokinase-1
334 and 2 (Mathupala and others 2001), and pyruvate dehydrogenase kinase (PDK)-1
335 (Kim and others 2006; Papandreou and others 2006) and -3 (Denko and others
336 2003) are just some in the long list of glycolytic enzymes that have been shown to
337 have increased expression in response to hypoxia and HIF activation. In normal
338 conditions, glucose transporters import glucose into the cytosol where it undergoes
339 glycolytic metabolism to produce pyruvate. Pyruvate can then be converted to
340 lactate or transported into the mitochondria. In the presence of oxygen, most cells
341 further degrade pyruvate through PDC, the committed step providing entry into the
342 TCA cycle. The cell may then further catabolize pyruvate to enable oxidative
343 phosphorylation and increased energy production. Full oxidation of carbons from
344 glucose through oxidative phosphorylation generates a 17-fold increase in
345 adenosine tri-phosphate (ATP) levels compared to anaerobic pathways. In 1857
346 Louis Pasteur recognized that in the presence of oxygen, cells metabolize glucose
347 via cellular respiration in the mitochondria; however, in hypoxic environments, cells
348 undergo fermentation. This is called the “Pasteur Effect” [(Pasteur 1857; Pasteur
349 1859) (reviewed in English (Racker 1974))]. Recent understandings of molecular
350 biology and biochemistry indicate that in response to HIF activation, PDK is
351 upregulated. PDK inhibits PDC activity by phosphorylation at three serine sites. PDC
352 inhibition drives fermentation, the phenomenon identified over a century and a half
353 ago. HIF activation and alterations to glucose metabolism have subsequently been
354 shown in many pathologic states, including cancer.

HIF and the Warburg Effect

In 1924 Otto Warburg observed that cancer cells preferentially engaged glycolytic metabolism and terminated at the conversion of pyruvate to lactate (Warburg 1930; Warburg 1956). This occurred in the presence of oxygen, the opposite of the Pasteur Effect previously discussed where normal cells metabolize glucose through the TCA cycle and promote mitochondrial respiration under these conditions. The production of lactate via glycolysis in normoxic conditions has since been dubbed the “Warburg Effect”, also known as “aerobic glycolysis” in the literature, and is less efficient at producing ATP than complete metabolism through the TCA cycle and oxidative phosphorylation. This metabolic phenotype is commonly seen in cancer cells, as well as HIF activation in normoxic conditions (Kim and others 2007; Koukourakis and others 2005; McFate and others 2008; Robey and others 2005; Semenza 2007).

Our lab has recently shown that decreasing the lactate production by knocking down PDK translation via constitutively expressed small inhibitory hairpin RNA (shRNA) causes a shift from aerobic glycolysis (the “Warburg Effect”) to full oxidation of glucose and results in decreased metastatic qualities in tumor cell lines (McFate and others 2008). Therefore manipulation of a key metabolic pathway may be able to change metastatic phenotypes in cancerous tissues, thus identifying possible new therapeutic targets.

Cancer hypermetabolism

A diagnosis of cancer is reserved for uncontrolled cell growth with metastatic phenotypes like migration, invasion and metastasis into surrounding tissues

(Hanahan and Weinberg 2000). CNS cancers are among the most invasive and deadly observed. One type of CNS cancer known as glioblastoma multiforme (GBM) is an astrocyte-derived grade 4 cancer that results in death on average from six to 12 months following diagnosis (Burton and Prados 2000; Miller and Perry 2007; Visted and others 2003). GBMs are notoriously resistant to therapeutic intervention and, due to the compacted and highly specific organization of the brain, are difficult to fully extract through surgical procedures (Burton and Prados 2000; Miller and Perry 2007; Nieder and others 2005; Terzis and others 2006). GBMs do metastasize and become increasingly deadly, though this metastatic event is rarely observed (Medhkour and Chan 2005; Mujic and others 2006; Newton and others 1992; Saad and others 2007). The aggressive nature of GBMs also makes it difficult to identify and treat these diseased tissues before the cancer becomes established or metastasizes (Aldape and others 2003; Nieder and others 2005; Terzis and others 2006).

The metastatic, metabolic, and proliferative qualities of GBMs exert a high demand of resources on surrounding CNS tissues. These qualities also cause healthy tissue to become necrotic, engulfed and degraded, or compacted leading to dysfunction (Burton and Prados 2000; Miller and Perry 2007). These cancerous phenotypes are also studied *in vitro* through cell culture and xenograph studies to better understand the basic biochemistry and cellular biology of GBMs. One such experimental model is the C6 glioma rat cell line, derived in the late 1960s by exposure of rat astrocytes to N,N'-nitroso-methylurea (Benda and others 1968). In a review article of *in vitro* experimental models of cancer, data on C6 glioma cells were

compiled and analyzed as a model closely associated to GBMs with respect to:
morphology; adhesion protein expression and activity; metastatic behaviors; and
biochemistry (Grobbs and others 2002). It is for these reasons that the studies
described in this work uses genetic transformations in C6 cells.

Branched-chain amino acids in cancer

Cancerous tissues are in a state of hypermetabolism due to increased
catabolic and anabolic demands for producing double the amounts of energy,
protein, and lipids for daughter cells. While glucose metabolism is altered in
cancerous tissues [the theme of my colleague's work (McFate and others 2008)],
other essential metabolic and physiologic pathways are also dysregulated in
cancerous diseases. Branched-chain amino acids (BCAAs) are a subset of essential
amino acids, categorized as such due to human inability to synthesize these basic
building blocks *de novo*. Given their multipotent roles in catabolism, anaplerosis,
anabolism, and signaling, it may be no surprise that a disease such as cancer can
have direct and indirect effects on BCAA pathways.

Free amino acids circulate throughout the body and also accumulate in small
pools in the intracellular environment. Hypermetabolic cancerous tissue can deplete
the human body of these essential amino acids. Although BCAAs are involved in
many pathways, both catabolic and anabolic, studies have shown that tumor-bearing
physiology drives hypoanabolism and hypercatabolism of the skeletal muscle in
animal models (Lorite and others 1997; Strelkov and others 1989). When this occurs
in humans with cancer, called cachexia, muscle wasting is often severe (Lundholm

and others 1976). Likewise, the breakdown of BCAA-rich skeletal muscle in the tumor-bearing state is documented in both the clinical setting (Hunter and others 1989; Inculet and others 1987; Inui 2002; O'Keefe and others 1990) and animal models (Baracos and Mackenzie 2006; Tessitore and others 1993; Whitehouse and others 2001). While the breakdown of skeletal muscle elevates BCAA levels for use in catabolism, the general health of the organism is impaired making resilience and recovery to therapy immensely difficult.

The tumor-bearing state is taxing to the organism, and metabolism is elevated in cancers. BCAAs are estimated to contribute 2-5% of total energy production in normal tissues, while they compose almost 25% of new proteins (Harper 1989). The precise pathways utilized in the cancerous or tumor-bearing state are unclear. However, where cachexia is a resultant influence of the cancerous tissue on the organism, the cancerous tissue itself requires the use of BCAAs for catabolic, anaplerotic, and anabolic pathways. The additional necessity of anabolism in these tissues makes them our primary target for metabolic manipulation. The proliferative and invasive phenotypes of some cancers require that BCAAs are supplied for the production of new proteins to build daughter cells upon division.

Branched-chain Amino Acids

BCAAs are important to many physiologic processes and have unique qualities, particularly to the human metabolic and molecular biological systems. They belong to a group of amino acids called essential amino acids that cannot be

synthesized in humans², which indicates that they must be obtained by dietary consumption. Once introduced into the physiological system of humans, the essential amino acids are shuttled and shunted to various pathways; the BCAAs are no different. Due to recent discoveries, it may be beneficial to make a distinction between the classical and non-classical understandings of BCAAs where the classical refers to metabolic pathways and the non-classical refers to signaling pathways.

Metabolism through the BCKDC

BCAAs largely bypass hepatic metabolism and are circulated throughout the body. Once BCAAs are within the cell membrane, they can proceed through many different metabolic fates; however, they must first be deaminated and/or decarboxylated in order to enter the biochemical system. Amine groups may be transferred between the carbon skeletons of amino acids by a number of aminotransferases that usually operate at or near equilibrium. There are two isotypes of branched-chain amino acid transaminases (BCAT); one is localized to mitochondria (BCATm) and is widely expressed in most tissues, and one is localized to the cytosol (BCATc) and selectively expressed in the brain, ovary, and placenta (Bixel and others 1997; Hall and others 1993; Hutson and Hall 1993; Hutson and others 1992; Ogawa and others 1970). The kinetic constants for both species are similar, with the K_M values for leucine (1–1.3mmole/L) and α -ketoglutarate (~0.6 mmole/L) exceeding the endogenous brain concentration of either substrate (~0.2 mmole/L) (Erecinska and others 1984). Deamination liberates an amine group that

² other essential amino acids are lysine, methionine, phenylalanine, threonine and tryptophan

can be used in classical processes, such as urea synthesis, and non-classical processes, which are discussed in detail later. The resulting deaminated BCAA is a carboxylic acid, branched-chain α -keto acid (BCKA). The BCAT enzyme operates on LeChatlier's principle of equilibrium since deamination is an easily reversible process. As such, the decarboxylation of the BCKAs is the most important step for their catabolism in humans (reviewed in (Harper 1989; Harper and others 1984)).

Decarboxylation occurs through the BCKDC. This complex belongs to a family of three dehydrogenase complexes including pyruvate dehydrogenase complex (PDC) and 2-oxo-glutarate dehydrogenase complex. These dehydrogenase complexes share the same basic structure, perform the same basic reactions, and all require the same set of cofactors: thiamine pyrophosphate, FAD, NAD, lipoate, and coenzyme A (CoA). The BCKDC is organized around a cubic core consisting of 24 lipoate-bearing dihydrolipoyl transacylase (E2) subunits, associated with the branched-chain α -keto acid decarboxylase/dehydrogenase (E1), dihydrolipoamide dehydrogenase (E3), BCKDC kinase (BDK), and BCKDC phosphatase (BDP). The E1 is formed by the α and β proteins interacting to form the two most important structures of the complex, the binding pocket for BCKAs and a crucial exposed Ser293 residue. Phosphorylation status of Ser293 indicates the activity state of BCKDC and is targeted by a specific kinase (Popov and others 1992; Shimomura and others 1990) and a phosphatase (Damuni and others 1984; Damuni and Reed 1987), although recent evidence has suggested isolation of another phosphatase (Joshi and others 2007). It has been established that the most potent inhibitor of BDK is the endogenous transamination product of leucine, KIC, indicating

that these metabolites promote their own metabolism. Decarboxylation of the BCKAs is the committed step in the catabolism of these molecules. After processing through the BCKDC, the metabolic products can be converted to acetyl-CoA and degraded for energy production (catabolism), used for fatty acid synthesis and energy storage (anabolism), or converted into intermediates to sustain biochemical pathways (anaplerosis). The combination of BCAA transport, BCAT reversible deamination of BCAAs, and the decarboxylation of BCKAs through BCKDC justify the supposition that BCKDC activity and intracellular concentrations of BCAAs and BCKAs govern the direction of BCAT activity.

Complete catabolism of BCAAs and BCKAs occurs only part of the time. Once BCKAs have proceeded through the BCKDC, they result in ubiquitously produced small molecules involved in various metabolic pathways, such as gluconeogenesis, the citric acid cycle, and fatty acid synthesis. However, BCAAs and BCKAs are involved in alternate pathways prior to catabolism through BCKDC.

New signaling roles have emerged (the comprehensive model)

In contrast to the classical understanding of BCAAs as energy substrates, new discoveries have made way for an emergent non-classical signaling paradigm. A few notable examples occur in the CNS where BCAAs are transported across the blood-brain barrier by the LAT1 and 2 transporters. Once inside the neurochemical milieu, BCAAs are used for glutamate homeostasis. One-third of all amine groups incorporated into glutamate, the brain's primary excitatory neurotransmitter, were found to originate in BCAAs; leucine alone supplies 30-50% to both glutamate and

glutamine, an important glutamate precursor and biochemical shuttling molecule (Kanamori and others 1998; Yudkoff and others 1990; Yudkoff and others 1983).

Although BCAA biochemistry is extensive, in both its nature and the study of the different pathways, there are also signaling pathways involving the BCAAs. The BCAAs leucine and valine are known to play important roles in the regulation of protein synthesis through interactions with translational initiation factors, including the mammalian target of rapamycin (mTOR), although leucine has been shown to be the most potent in stimulating the mTOR pathway (Figure 1) (Anthony and others 2000a; Kimball and Jefferson 2006; Lynch and others 2003). mTOR is a global regulator of cellular and molecular processes concerning cell survival. Briefly, activation of mTOR, a protein kinase, begins a cascade of events promote protein translation, such as: phosphorylation of 4E-binding protein 1 (4E-BP1) for the release of eukaryotic initiation factor-4E (eIF-4E) (Brunn and others 1997; Kimball and others 1996; Xu and others 1998); phosphorylation of S6 ribosomal protein kinase (S6RK) to activate downstream targets (Kimball and others 1999; Long and others 2000); and ribosomal biogenesis, although this is still considered a hypothesis by many which is refuted by some (Stolovich and others 2002; Tang and others 2001).

The most recent data detailing leucine's non-classical signaling role has been presented which details a further role of leucine-mTOR signaling in the hypothalamus (Cota and others 2006). Cota *et al* (2006) reported that leucine, acting through mTOR and other translational regulators, was able to initiate signaling to

regulate fuel availability and usage via the arcuate and paraventricular nuclei in the rat brain. These nuclei are located near the medial eminence, a circumventricular organ in the brain with a direct “window” to the blood supply. Such gaps in the blood-brain barrier allow the careful monitoring and regulation of metabolites and hormones. As a result, hypothalamic signaling can occur adjusted to dynamic systemic needs based on feeding and fasting. The authors found that central administration of leucine resulted in increased hypothalamic mTOR signaling and decreased food intake and body weight (Cota and others 2006). This study further elaborated on leucine’s importance as a metabolic signaling molecule.

Summary and Segue

In light of the previously defined complex nature of BCAAs in physiological systems, we examined the role of BCAAs in HIF biology and cancer biochemistry. Our research questions were: what is the effect of increased BCAA levels in an organism with cancer (seen in rats and humans); and is there a greater role for BCAAs in cancer metabolism, especially in light of the fact that PDC is very clearly a metabolic switch? As previously stated, BCAA transaminated products (BCKAs) have been shown to stabilize HIF at high *in vitro* levels. Our first question is concerned with what occurs when BCAAs and BCKAs are present at lower levels and if HIF activation is robust enough to drive HIF-targeted transcriptional upregulation. Our second question is concerned with the hypermetabolic state present in cancer cells and if BCKDC can be manipulated in order to cause significant changes to cancer cell phenotypes.

Branched-chain α -ketoacids increase HIF-1 signaling independently of mTOR

Jeremy HENRIQUES^{*1} and Ajay VERMA, MD PhD^{*†}

^{*}Neuroscience Program, Uniformed Services University of the Health Sciences,
Bethesda, MD

[†]Department of Neurology, Uniformed Services University of the Health Sciences,
Bethesda, MD

¹To whom correspondence should be addressed

Jeremy Henriques

Neuroscience Program

Uniformed Services University of the Health Sciences

4301 Jones Bridge Road

Bethesda, MD 20814

Telephone: (301) 295-3840

Fax: (301) 295-3825

e-mail: henriques.jeremy@gmail.com

Abbreviations: BCAA, branched-chain amino acid; BCKA, branched-chain α -

ketoacid; BCKDC, branched-chain α -ketoacid dehydrogenase complex; BCKDH,

branched-chain α -ketoacid dehydrogenase; BDK, BCKDH kinase; GFP, green

fluorescent protein; FIH, factor inhibiting HIF; HIF, hypoxia inducible factor; KIC, 2-

ketoisocaproate; KIV, 2-ketoisovalerate; KMV, 2-keto-3-methyl valerate; mTOR,

mammalian target of rapamycin; ODD, oxygen-dependent degradation domain;

PDH, pyruvate dehydrogenase; PHD, prolyl hydroxylase domain proteins; S6R, S6 ribosomal protein; VEGF, vascular endothelial growth factor

Summary

Branched-chain amino acids and ketoacids participate in several biochemical pathways and are emerging as novel signaling molecules. Leucine activates the mammalian target of rapamycin (mTOR) kinase, while its deaminated metabolite alpha-ketoisocaproic acid (KIC) can stabilize the hypoxia-inducible transcription factor, HIF-1. Since mTOR has been implicated in HIF-1 stabilization, we investigated whether the ability of KIC to stabilize HIF-1 involved mTOR activity. In rat C6 glioma cells, KIC treatment promoted the accumulation of HIF-1 α as well as a stably transfected green fluorescent protein containing the HIF-1 α oxygen-dependent degradation domain. Inhibition of the HIF-1 decay mechanism by KIC was correlated with increased VEGF secretion by C6 cells. KIC also increased phosphorylation of the mTOR target protein S6R. However, while rapamycin treatment inhibited KIC-induced S6R phosphorylation, it did not affect KIC-induced HIF-1 stabilization or VEGF secretion. Instead, these rapamycin-insensitive KIC effects were selectively reversed by ascorbate, a cofactor required by the proly-hydroxylase domain proteins, which control HIF-1 decay. KIC also promoted ascorbate sensitive HIF-1 α accumulation and VEGF secretion in primary rat astrocytes. These results suggest that KIC may reversibly inactivate proly-hydroxylase domain proteins independently of mTOR activity.

Introduction

The branched-chain amino acids (BCAAs: leucine, isoleucine, and valine), whose side chains contain a branched methyl group, are essential amino acids known to be involved in several biochemical pathways. Leucine, isoleucine, and valine are deaminated by a specific branched chain amino acid transferase to their respective branched chain α -ketoacids (BCKAs) 2-ketoisocaproate (KIC), 2-ketoisovalerate (KIV), and 2-keto-3-methylvalerate (KMV) (Hutson and others 1988). BCKAs are then catabolized through the mitochondrial branched-chain α -keto acid dehydrogenase complex (BCKDC), which is composed of multiple subunits, including the regulatory branched chain ketoacid dehydrogenase E1- α subunit (BCKDH- α) (Danner and others 1978; Parker and Randle 1978; Pettit and others 1978; Roberts and Sokatch 1978). BCKDC is regulated by reversible inhibitory phosphorylation of BCKDH- α through the actions of a branched chain dehydrogenase kinase (BDK) (Popov and others 1992; Shimomura and others 1990) and an unidentified phosphatase (Damuni and others 1984; Damuni and Reed 1987). The catabolism of BCAAs via the transaminase and the BCKDC generates acyl-CoA intermediates which undergo dehydrogenation. Ultimately, leucine is converted to acetyl-CoA and acetoacetate; isoleucine to acetyl-CoA and succinyl-CoA; and valine to succinyl-CoA. In some tissues, these final products can be fully oxidized via the citric acid cycle, while in others these are directed toward the synthesis of ketone bodies (acetoacetate and acetyl-CoA) and glucose (succinyl-CoA) (Greenberg and Reaven 1966; Noda and Ichihara 1974; Noda and Ichihara 1976).

BCAAs are also used as anabolic building blocks for de novo protein synthesis. Whereas all three BCAAs serve equally prominent roles in these biochemical pathways, leucine has emerged as an important signaling molecule as well. Leucine has been shown to stimulate protein synthesis via mobilization of the protein translation machinery through activation of mammalian target for drug rapamycin (mTOR) activity (Kimball and others 1999). Leucine also promotes insulin synthesis and secretion (Lambert and others 1986), and inhibits autophagy (Mordier and others 2000). In biochemical pathways restricted to central nervous system tissue leucine plays an important role in maintaining homeostasis of glutamate, the brain's major excitatory and most abundant neurotransmitter (Yudkoff and others 1996; Yudkoff and others 1990; Yudkoff and others 1983). More recently, signaling actions of leucine in the hypothalamus have been implicated in the regulation of feeding behavior (Cota and others 2006).

Accumulation of BCAAs and BCKAs has also long been known to contribute to the pathogenesis of BCKDC deficiency, which is commonly known as maple syrup urine disease (Dancis and others 1960). Symptoms of this condition begin in early infancy and include poor feeding, vomiting, dehydration, lethargy, hypotonia, seizures, ketoacidosis, and neurological decline (Chuang and others 2006). However, the pathophysiology underlying the nervous system effects remains unclear and has not been linked to the signaling roles identified for BCAAs. A better appreciation of BCAA signaling roles may thus help clarify the regulation of important physiological and pathological processes.

645 Novel signaling actions of many other metabolic intermediates have also
646 become elucidated in recent years. Glycolytic metabolites and tricarboxylic acid
647 (TCA) cycle intermediates have also been shown to promote insulin secretion
648 (MacDonald and others 1989), regulate hypothalamic hunger signals (Lam and
649 others 2005), induce angiogenesis (Murray and Wilson 2001), and promote
650 stabilization of the hypoxia inducible transcription factor HIF-1 (Isaacs and others
651 2005; Lu and others 2005; Lu and others 2002; Pollard and others 2005; Selak and
652 others 2005). Cell survival in hypoxic environments is critically dependent upon HIF-
653 1 in both normal and neoplastic tissues (Dalgard and others 2004; Lu and others
654 2005; Lu and others 2002). Our recent study evaluating the effect of BCAAs on HIF-
655 1 α stabilization showed that BCKAs could also stabilize HIF-1 α levels (Lu and others
656 2005).

657 Under normal oxygen conditions (or normoxia) HIF- α is constitutively
658 synthesized but rapidly degraded by specific HIF prolyl hydroxylases, referred to as
659 prolyl hydroxylase domain proteins 1-3 (PHD 1-3) (Wang and others 1995). The
660 PHD enzymes require the cofactors iron, ascorbate (Knowles and others 2003) and
661 the tricarboxylic acid [TCA] cycle intermediate α -ketoglutarate (Bruick and McKnight
662 2001; Ivan and others 2001; Jaakkola and others 2001), for sustained enzymatic
663 activity. PHDs transfer hydroxyl (-OH) groups derived from dissolved O₂ onto two
664 proline residues located in the oxygen-dependent degradation (ODD) domain of the
665 HIF- α protein (Hon and others 2002; Masson and others 2001; Min and others 2002;
666 Srinivas and others 1999). This oxygen dependent post-translational modification
667 acts as a recognition signal for the ubiquitin-mediated degradation of HIF- α subunits.

In the absence of atmospheric oxygen, HIF- α protein levels increase, dimerize with HIF- β (Gradin and others 1996; Kallio and others 1997; Wood and others 1996), translocate to the nucleus, and activate transcription of many genes that promote hypoxic survival. Another HIF- α hydroxylase known as the factor inhibiting HIF (FIH) governs the hydroxylation of an asparagine residue on HIF- α (Hewitson and others 2002; Lando and others 2002a; Lando and others 2002b; Sang and others 2002), thus regulating its association with other transcriptional cofactors.

Along with hypoxia, PHD enzyme activity can be inhibited by reducing the interaction of cofactors with the enzyme. Thus, iron chelators, α -ketoglutarate analogues (Bruick and McKnight 2001; Ivan and others 2001; Jaakkola and others 2001), and ascorbate deficiency (Knowles and others 2003) can blunt PHD activity and enhance HIF-1 accumulation. In fact, the effect of on BCKAs on HIF-1 stabilization was discovered through a screen of biological α -ketoacids with structural similarity to α -ketoglutarate (Lu and others 2005). However, given the wider emerging signaling roles of BCAAs, it is possible that the BCKA effect we observed on HIF-1 also involved other pathways. In particular, mTOR has been shown to be both an upstream regulator of HIF signaling in cancer cells (Hudson and others 2002) and a regulatory target potentially influenced by leucine (Anthony and others 2000b). Thus it remains unknown whether the BCKA effect on HIF-1 regulation is mediated selectively via the PHDs or via mTOR. Given the potential widespread impact of α -ketoacid signaling mechanisms, the present study further investigated the role of BCKAs (and BCAAs) in the stabilization of HIF- α to clarify this question.

691

692 **Experimental Methods**

693 All chemicals were purchased from Sigma-Aldrich and cell culture products were
694 purchased from GIBCO, unless otherwise stated.

695 **Transgenic ODD-GFP C6 Glioma Cell Line**

696 The generation of the ODD-GFP C6 glioma cell line has been previously
697 described (D'Angelo and others 2003). The ODD is the portion of the HIF-1 α protein
698 that is hydroxylated and subsequently degraded by the ubiquitin ligase molecular
699 machinery. Fusion of the HIF-1 α ODD to a GFP molecule allowed GFP
700 immunodetection to serve as a probe for the PHD mediated degradation of HIF-1 α .

701 **Cell culture and Chemical Treatments**

702 Cells were cultured in Dulbecco's Modified Eagle Media (DMEM, Invitrogen)
703 supplemented with 10% fetal bovine serum (FBS), 1% penicillin-streptomycin, and
704 1.5 mg/mL G418 (Gibco). G418 was used to select transgenically altered cells. Cell
705 lines were maintained in 21% O₂, 5% CO₂, and 74% N₂ in a humidified cell incubator
706 at 37°C. Chemical treatments were performed in Krebs Saline Buffer and incubation
707 times are as indicated. Each chemical treatment was performed by dilution from
708 100X stock in Krebs saline buffer. For cell hypoxia treatment, the culture dishes
709 were sealed in a modular incubator chamber, flushed with gas containing 1% O₂,
710 5%CO₂, and 94% N₂ for 5 min, and incubated in this environment at 37°C for the
711 indicated times.

712 **Primary Astrocyte-enriched Cell Cultures**

Primary cultures from derived from rat postnatal day 2 cerebral cortex were prepared as described (Armstrong 1998). Briefly, brains were surgically extracted, digested with protease, and plated in poly-D-lysine coated tissue culture flasks. Cultures were maintained in DMEM with 10% fetal bovine serum (FBS): DMEM (Life Technologies) supplemented with 1 mM sodium pyruvate (Sigma) and 25 mg/ml gentamicin (Life Technologies). Following 10 days in culture the flasks were placed on a rotary shaker (190rpm) for 18h to dislodge immature oligodendrocyte lineage cells and microglia. The remaining cells were astrocyte-enriched populations. Astrocyte cultures were refreshed every 72h with DMEM medium containing 10% FBS supplemented with 1mM sodium pyruvate and 25mg/ml gentamicin.

Western Blot and Densitometry Analyses

Cells were washed three times with cold PBS. Appropriate amount of lysis buffer containing Radio Immuno Precipitant Assay (RIPA) buffer (Tris-HCL, pH 7.4 [50mM], NaCl [150mM], NP-40 [1%], Sodium deoxycholate [0.5%], SDS [0.1%], and EDTA [5mM]; Bioworld), 1% SDS, and 1X protease inhibitor cocktail (Roche) then scraped. Cell remnants were then collected in 1.5 ml microcentrifuge tubes. The cell material was sonicated for 25 seconds at 50 Hz, then placed in rack at room temperature. Cell lysates were then spun for 5 minutes at 12,000 g and the supernatants were transferred to fresh tubes. Protein levels were determined by the BCA method of analysis (Pierce). Proteins were separated on 4-12% Bis-Tris SDS-polyacrylamide gradient gels (Invitrogen) and transferred to nitrocellulose membranes (Invitrogen). Membranes were blocked using 5% goat or horse serum (Vecotr Labs) in Tris buffered saline with 0.1% Tween-20. Antibodies used were

anti-HIF-1 α 1:500 (Biossource, Novus), anti-HIF-1 β 1:1000 (Biossource, Novus), anti-phospho-Ser292-E1 α BCKDH (generous gift from C. Lynch, 1:20,000) (Lynch and others 2003), anti-E1 α BCKDH (generous gift from C. Lynch, 1:1000) (Lynch and others 2003), anti-GFP 1:1000 (Roche), anti-phospho-S6R 1:1000 (Cell Signaling), and anti- β actin 1:10,000 (AbCam). Protein bands were visualized by enhanced chemiluminescence (Pierce) using either Kodak film and developer or CCD luminescence camera (Fuji Film). Signals were quantified using densitometry in ImageJ (Wayne Rasband, NIH). Phosphoproteins and HIF-1 α were normalized to total protein amounts and HIF-1 β , respectively, unless otherwise stated. The β -actin levels of non-phosphoproteins were statistically compared. If β -actin levels differed, the data were not used in the subsequent analysis. Graphs indicate arbitrary units on the y-axes derived from signals normalized to non-treated or wild type controls by direct comparative ratios. These normalizations were necessary to control for protein loading.

Enzyme Linked Immunosorbent Assay

Cells were cultured to 100% confluency in 6-well cell culture plates. Experimental conditions were applied and the cells were incubated for 12 hours under culture conditions. 50 μ L of media was extracted and used in a Quantikine Rat VEGF (vascular endothelial growth factor) ELISA assay kit (R&D Systems Inc.) Absorbance was read at 405nm generating pg/mL VEGF protein. Cells were trypsinized and counted using the ViCell Trypan Blue automated cell counter subsequent to media extraction for ELISA. Cell numbers were used to normalize the

data to account for differential plating and proliferation confounders. Data is represented as pg VEGF protein per mL media per hundred cells.

Cell counting

Cell were washed and trypsinized for cell counting procedures for ELISA normalization. Trypsin was inactivated by adding equal amounts of media with 10% FBS. Five hundred μ L of suspensions of each preparation were then placed in a Vi-Cell Trypan Blue automated counter for cell counting.

Statistical Analyses

Statistical analyses ANOVA, Tukey's HSD and Fisher's PLSD post hoc tests were performed using STATview statistical software. Where it was necessary to compare treatment only to control groups, ANOVA and Dunnett's post hoc analyses were performed in SPSS statistical software.

Results

KIC induced HIF-1 stabilization is independent of mTOR activity

KIC can be reversibly transaminated to produce leucine, which can activate mTOR, a kinase that regulates protein translation via phosphorylation of proteins such as S6R. In some studies, mTOR activation has been implicated in increasing HIF-1 α (Hudson and others 2002) and VEGF protein levels (Humar and others 2002), suggesting that this could be one mechanism for the previously observed HIF-1 α stabilization by KIC. Alternatively, stabilization of HIF-1 α by α -ketoacids has been proposed to result from a reversible inactivation of PHD enzymes (Lu and others 2005). This conclusion is supported by the rapid reversibility of α -ketoacid mediated HIF-1 α accumulation by ascorbate and Fe²⁺ (Lu and others 2005; Lu and

others 2002). To determine which of these mechanisms is involved in KIC induced HIF-1 α accumulation we sought to determine the relative sensitivity of the KIC-induced HIF-1 α accumulation to either ascorbate or rapamycin, a well known mTOR inhibitor (Brown and others 1994; Sabatini and others 1994; Sabers and others 1995). We employed immunodetection of pSer^{235/236}S6R to assess mTOR activity and ODD-GFP stabilization to assess PHD activity.

As shown in figure 1A and 1B, when ODD-GFP transfected C6 glioma cells were treated with hypoxia (1% O₂) vs. KIC (2mM) for 3h, only KIC treatment demonstrated significant increase of pSer^{235/236}S6R immunoreactivity in cell extracts. Moreover, rapamycin treatment (0.1ng/ml) completely prevented S6R phosphorylation while ascorbate treatment (100 μ M) did not. In the same cell extracts, we also examined changes in HIF-1 α and HIF-1 β immunoreactivity. As shown in figure 1A and 1C, both hypoxia and KIC treatment led to a significant accumulation of HIF-1 α with no change in HIF-1 β . The KIC induced HIF-1 α accumulation was clearly far more sensitive to inhibition by ascorbate than by rapamycin. ODD-GFP accumulation was also stimulated by KIC and this effect was selectively inhibited by ascorbate but not by rapamycin (figure 1A, 1D).

Functional HIF-1 induced gene expression requires nuclear translocation of the HIF-1 α /HIF-1 β heterodimer and interaction with the transcriptional coactivator CBP/P300 (Arany and others 1996). This latter step is normally kept inhibited by the asparagine hydroxylase FIH (Lando and others 2002a; Lando and others 2002b; Sang and others 2002). Full gene expression also requires DNA transcription and mRNA translation to produce functional protein. In order to assess whether the KIC

induced HIF-1 α accumulation that we observed was effective in activating gene expression, we measured the media accumulation of VEGF protein, a well known target of HIF-1 α following 12h culture. As shown in figure 1E, both hypoxia and KIC significantly increased media VEGF levels and the KIC effect was selectively blunted by ascorbate and not by rapamycin. Similar results were also seen with leucine treatment (figure 1F). These results supported the notion that KIC induced HIF-1 α accumulation is mediated by reversible inactivation of PHD activity and is not significantly impacted by mTOR activation.

HIF-1 α stabilization and BCKD α phosphorylation show reciprocal dose responses to KIC

The BCKAs KIC, KIV, and KLV have been shown to increase HIF-1 α levels in normoxic conditions at high doses. At these high doses these α -ketoacids may have antioxidant and other non-specific actions. Moreover, in living cells, these ketoacids may also be rapidly metabolized through BCKDC activity. BCKDC normally regulates the BCKDC through inhibitory phosphorylation of the BCKDH α subunit at Ser292. KIC is the chief endogenous inhibitor of BCKDC, a mechanism that normally allows feed-forward activation of the BCKDC to maintain low levels of KIC. These complex fuel routing mechanisms may thus impact the selectivity of signaling functions for KIC. We therefore determined whether the state of inhibitory BCKDH α phosphorylation at Ser292, correlated with the effect of KIC on HIF-1 α stabilization. To accomplish this, C6-ODD-GFP cells were treated with concentrations of KIC ranging from 1 to 3000 μ M. The treated cells were then harvested for analysis of HIF-1 α levels and phosphorylation status of BCKDH- α Ser292. A representative

dose-response curve for KIC effect on HIF-1 α accumulation and BCKDH- α pSer292 levels via Western blotting is shown in Fig. 2A. Densitometric analysis of blots is shown in Fig. 2B with standardization to non-treated controls. Increased HIF-1 α signal relative to non-treated controls could be seen at the 10 μ M dose of KIC ($p=0.001$) and all concentrations from 30 μ M KIC to 3000 μ M KIC showed significant HIF-1 α accumulation ($p<0.0001$). Phosphorylation of BCKDH- α Ser292 is shown as the ratio of phosphorylated to total protein for internal standardization. Treatments of cells with 1, 10 and 30 μ M KIC resulted in significantly decreased phosphorylation ratios of BCKDH- α with respect to non-treated controls ($p<0.0001$) with no BCKDH- α phosphorylation being observed at higher doses. These data indicated that despite inhibition of BCKDH phosphorylation, BCKAs such as KIC can still promote HIF-1 stabilization. We did however observe a complex dose response curve for KIC in stabilizing HIF-1 which suggested the interaction of more than one process to the overall effect.

KIC and leucine induced VEGF production displays a biphasic response

Given our demonstration of the ability of KIC to participate in distinct signaling and metabolic pathways in the same cell type, we wondered whether the overall production of HIF-1 mediated gene expression by this α -ketoacid would show a linear dose response or one that was more complex. We therefore investigated the dose response of KIC in promoting an increase in media VEGF. C6-ODD-GFP cells were treated with increasing concentrations of KIC as well as leucine ranging from 100 to 5000 μ M. Positive and negative controls were also included using hypoxia (1% O₂) and vehicle-treated conditions, respectively. The treated cell media was

then harvested after 12h and analyzed by the ELISA technique. Cells were then counted with the ViCell Trypan blue automated cell counter. We found increased secretion of VEGF protein in hypoxia positive-control condition (Figs. 2C and D; $p < 0.0001$). However, the dose-response curves for KIC and leucine were bimodal (Fig. 2C and D). Thus while 0.1mM KIC produced significant increase in VEGF production, 0.5mM KIC did not. However, subsequently increasing concentrations of KIC did produce a significant increase in VEGF. A similar biphasic dose response was observed with leucine although the required effective dose range was higher than that for KIC.

KIC stabilizes HIF-1 α and promotes VEGF expression in primary astrocyte cultures

We also addressed whether the effects of KIC that we observed in the C6 glioma cell line could be observed in primary cultures of non-transformed cells. We thus treated astrocyte-enriched primary rat brain cultures with hypoxia, or with KIC [0.1 and 1.0 mM] with and without ascorbate [100 μ M]. Media from treated cells were analyzed by ELISA (Fig. 3C). Treated cells were harvested and analyzed by Western blotting for HIF-1 α and - β . A representative Western blot of analyses is shown in Fig. 3A. Densitometry of these blots is plotted in Fig. 3B. The positive-control hypoxia treatments resulted in significant increases from all other treatments ($p < 0.001$ for all comparison). Both the 0.1 mM and 1.0 mM KIC treatments resulted in increased HIF-1 α/β ratios relative to non-treated controls ($p < 0.05$ for both comparisons). The results of the two KIC alone treatments did not differ statistically from one another. The KIC treatments showed significantly increased HIF-1 α/β

ratios vs. treatment with KIC and ascorbate ($p < 0.05$ for both comparisons). The two KIC treatment doses with ascorbic acid did not differ from each other. These data indicate that KIC indeed stabilizes HIF-1 α protein levels at normoxia in normal non-neoplastic cells as well.

KIC-induced, ascorbate-reversible VEGF protein expression was also observed in media taken from the primary astrocyte cultures (Fig. 3C). The non-treated controls showed much lower VEGF protein compared to the positive-control hypoxia ($p < 0.0001$) and the two doses of KIC ($p < 0.0001$). No differences were seen however between non-treated controls and either of the KIC treatments when ascorbic acid was included.

Discussion

BCAAs and BCKAs are known to be involved in multiple biochemical pathways and are also becoming appreciated for their signaling functions. Previous work from our lab identified BCKAs as being among a few naturally occurring α -ketoacids that were capable of stabilizing HIF-1 α levels (Lu and others 2005). Among the BCKAs this effect was greatest for KIC. In this report, we have elaborated on this previous work to show this action to be independent of mTOR. KIC was shown to activate signaling via mTOR as indicated by its enhancement of rapamycin sensitive S6 phosphorylation. KIC was also able to decrease phosphorylation of BCKDH α . However, the ability of KIC to promote HIF-1 α stabilization and produce VEGF elaboration appeared to result from an interference of PHD activity. This was supported by the ability of ascorbate to reverse both of these actions and also by the ability of KIC to promote ODD-GFP accumulation in an

896 ascorbate-reversible manner. The latter tool is driven by constitutive expression
897 under a CMV promoter and the protein product is degraded specifically by O₂-
898 dependent hydroxylation via the PHD enzymes (D'Angelo and others 2003).
899 Interestingly ascorbate treatment of C6 transgenic cells led to a decreased level of
900 ODD-GFP fusion protein as compared to non-treated controls (Figure 1D). This is
901 possibly due to the normal homeostatic relationship between ODD-GFP expression
902 and PHD activity which results in a basal level of protein (evidenced by the former)
903 and the addition of ascorbate may increase the PHD degradation cycle turnover
904 (evidenced by the latter), thus reducing the normal basal level to a non-detectable
905 signal.

906 In addition to these actions, BCKAs participate in transamination events,
907 possess antioxidant capabilities and also influence the bioenergetic status of cells.
908 Thus, it is not surprising that we observed unusual dose response curves for VEGF
909 elaboration by KIC and leucine. Although a clear explanation for this effect requires
910 further experiments, the differential dose dependent actions of KIC on the distinct
911 biological effects demonstrated and described above may account for this
912 phenomenon. It is possible that KIC and leucine are capable of inactivating PHDs at
913 low doses. However, BCKDC activation by higher doses of KIC due to inhibition of
914 BCKDH α phosphorylation may lower the effective concentration of KIC for
915 performing this action. Although other explanations may be possible, our results do
916 clearly demonstrate the ability of KIC to stabilize HIF-1 in living cells and show for
917 the first time the ability of KIC and leucine to promote VEGF production. This latter
918 action may have clinical relevance.

High accumulation of BCAAs and BCKAs is a hallmark of maple syrup urine disease (Dancis and others 1960). One of the most ominous presentations of this disease is cerebral edema (Brismar and others 1990; Riviello and others 1991). Our demonstration that KIC could induce HIF-1 α stabilization and VEGF elaboration in primary astrocytes suggests that this signaling mechanism may play a role in the pathogenesis of cerebral edema in maple syrup urine disease. This is because VEGF is well known to be a key contributor to edema through its action on vascular permeability (Josko and others 2000). Moreover, our demonstration that ascorbate can reverse KIC-induced VEGF elaboration from astrocytes suggests a possible simple treatment that can be tested in this clinical condition. Finally, given the prominent role of HIF-1 biology in cancer progression, our results suggest that a role for altered BCKDC biology in cancer should be further evaluated.

Acknowledgements: This work was funded by NIH grants NS73814 and CA113506 to AV.

References

- [1] S.M. Hutson, D. Fenstermacher, C. Mahar, Role of mitochondrial transamination in branched chain amino acid metabolism, J Biol Chem 263 (1988) 3618-3625.
- [2] D.J. Danner, S.K. Lemmon, L.J. Elsas, 2nd, Substrate specificity and stabilization by thiamine pyrophosphate of rat liver branched chain α -ketoacid dehydrogenase, Biochem Med 19 (1978) 27-38.

941 [3] P.J. Parker, P.J. Randle, Partial purification and properties of branched-chain
 942 2-oxo acid dehydrogenase of ox liver, *Biochem J* 171 (1978) 751-757.

943 [4] F.H. Pettit, S.J. Yeaman, L.J. Reed, Purification and characterization of
 944 branched chain alpha-keto acid dehydrogenase complex of bovine kidney, *Proc Natl*
 945 *Acad Sci U S A* 75 (1978) 4881-4885.

946 [5] C.M. Roberts, J.R. Sokatch, Branched chain amino acids as activators of
 947 branched chain ketoacid dehydrogenase, *Biochem Biophys Res Commun* 82 (1978)
 948 828-833.

949 [6] K.M. Popov, Y. Zhao, Y. Shimomura, M.J. Kuntz, R.A. Harris, Branched-chain
 950 alpha-ketoacid dehydrogenase kinase. Molecular cloning, expression, and sequence
 951 similarity with histidine protein kinases, *J Biol Chem* 267 (1992) 13127-13130.

952 [7] Y. Shimomura, N. Nanaumi, M. Suzuki, K.M. Popov, R.A. Harris, Purification
 953 and partial characterization of branched-chain alpha-ketoacid dehydrogenase kinase
 954 from rat liver and rat heart, *Arch Biochem Biophys* 283 (1990) 293-299.

955 [8] Z. Damuni, M.L. Merryfield, J.S. Humphreys, L.J. Reed, Purification and
 956 properties of branched-chain alpha-keto acid dehydrogenase phosphatase from
 957 bovine kidney, *Proc Natl Acad Sci U S A* 81 (1984) 4335-4338.

958 [9] Z. Damuni, L.J. Reed, Purification and properties of the catalytic subunit of
 959 the branched-chain alpha-keto acid dehydrogenase phosphatase from bovine kidney
 960 mitochondria, *J Biol Chem* 262 (1987) 5129-5132.

961 [10] R. Greenberg, G. Reaven, The effect of L-leucine on hepatic glucose
 962 formation, *Pediatrics* 37 (1966) 934-941.

- 963 [11] C. Noda, A. Ichihara, Control of ketogenesis from amino acids. II. Ketone
964 bodies formation from alpha-ketoisocaproate, the keto-analogue of leucine, by rat
965 liver mitochondria, J Biochem 76 (1974) 1123-1130.
- 966 [12] C. Noda, A. Ichihara, Control of ketogenesis from amino acids. IV. Tissue
967 specificity in oxidation of leucine, tyrosine, and lysine, J Biochem 80 (1976) 1159-
968 1164.
- 969 [13] S.R. Kimball, L.M. Shantz, R.L. Horetsky, L.S. Jefferson, Leucine regulates
970 translation of specific mRNAs in L6 myoblasts through mTOR-mediated changes in
971 availability of eIF4E and phosphorylation of ribosomal protein S6, J Biol Chem 274
972 (1999) 11647-11652.
- 973 [14] D.G. Lambert, K. Hughes, T.W. Atkins, Insulin release from a cloned hamster
974 B-cell line (HIT-T15). The effects of glucose, amino acids, sulphonylureas and
975 colchicine, Biochem Biophys Res Commun 140 (1986) 616-625.
- 976 [15] S. Mordier, C. Deval, D. Bechet, A. Tassa, M. Ferrara, Leucine limitation
977 induces autophagy and activation of lysosome-dependent proteolysis in C2C12
978 myotubes through a mammalian target of rapamycin-independent signaling pathway,
979 J Biol Chem 275 (2000) 29900-29906.
- 980 [16] M. Yudkoff, Y. Daikhin, L. Grunstein, I. Nissim, J. Stern, D. Pleasure,
981 Astrocyte leucine metabolism: significance of branched-chain amino acid
982 transamination, J Neurochem 66 (1996) 378-385.
- 983 [17] M. Yudkoff, I. Nissim, L. Hertz, Precursors of glutamic acid nitrogen in primary
984 neuronal cultures: studies with ¹⁵N, Neurochem Res 15 (1990) 1191-1196.

985 [18] M. Yudkoff, I. Nissim, S. Kim, D. Pleasure, K. Hummeler, S. Segal, [15N]
 986 leucine as a source of [15N] glutamate in organotypic cerebellar explants, *Biochem*
 987 *Biophys Res Commun* 115 (1983) 174-179.
 988 [19] D. Cota, K. Proulx, K.A. Smith, S.C. Kozma, G. Thomas, S.C. Woods, R.J.
 989 Seeley, Hypothalamic mTOR signaling regulates food intake, *Science* 312 (2006)
 990 927-930.
 991 [20] J. Dancis, M. Levitz, R.G. Westall, Maple syrup urine disease: branched-
 992 chain keto-aciduria, *Pediatrics* 25 (1960) 72-79.
 993 [21] D.T. Chuang, J.L. Chuang, R.M. Wynn, Lessons from genetic disorders of
 994 branched-chain amino acid metabolism, *J Nutr* 136 (2006) 243S-249S.
 995 [22] M.J. MacDonald, L.A. Fahien, R.J. Mertz, R.S. Rana, Effect of esters of
 996 succinic acid and other citric acid cycle intermediates on insulin release and inositol
 997 phosphate formation by pancreatic islets, *Arch Biochem Biophys* 269 (1989) 400-
 998 406.
 999 [23] T.K. Lam, R. Gutierrez-Juarez, A. Pocai, L. Rossetti, Regulation of blood
 1000 glucose by hypothalamic pyruvate metabolism, *Science* 309 (2005) 943-947.
 1001 [24] B. Murray, D.J. Wilson, A study of metabolites as intermediate effectors in
 1002 angiogenesis, *Angiogenesis* 4 (2001) 71-77.
 1003 [25] H. Lu, C.L. Dalgard, A. Mohyeldin, T. McFate, A.S. Tait, A. Verma, Reversible
 1004 inactivation of HIF-1 prolyl hydroxylases allows cell metabolism to control basal HIF-
 1005 1, *J Biol Chem* 280 (2005) 41928-41939.

1006 [26] H. Lu, R.A. Forbes, A. Verma, Hypoxia-inducible factor 1 activation by aerobic
 1007 glycolysis implicates the Warburg Effect in carcinogenesis, *J Biol Chem* 277 (2002)
 1008 23111-23115.

1009 [27] J.S. Isaacs, Y.J. Jung, D.R. Mole, S. Lee, C. Torres-Cabala, Y.L. Chung, M.
 1010 Merino, J. Trepel, B. Zbar, J. Toro, P.J. Ratcliffe, W.M. Linehan, L. Neckers, HIF
 1011 overexpression correlates with biallelic loss of fumarate hydratase in renal cancer:
 1012 novel role of fumarate in regulation of HIF stability, *Cancer Cell* 8 (2005) 143-153.

1013 [28] P.J. Pollard, J.J. Briere, N.A. Alam, J. Barwell, E. Barclay, N.C. Wortham, T.
 1014 Hunt, M. Mitchell, S. Olpin, S.J. Moat, I.P. Hargreaves, S.J. Heales, Y.L. Chung, J.R.
 1015 Griffiths, A. Dalglish, J.A. McGrath, M.J. Gleeson, S.V. Hodgson, R. Poulson, P.
 1016 Rustin, I.P. Tomlinson, Accumulation of Krebs cycle intermediates and over-
 1017 expression of HIF1alpha in tumours which result from germline FH and SDH
 1018 mutations, *Hum Mol Genet* 14 (2005) 2231-2239.

1019 [29] M.A. Selak, S.M. Armour, E.D. MacKenzie, H. Boulahbel, D.G. Watson, K.D.
 1020 Mansfield, Y. Pan, M.C. Simon, C.B. Thompson, E. Gottlieb, Succinate links TCA
 1021 cycle dysfunction to oncogenesis by inhibiting HIF-alpha prolyl hydroxylase, *Cancer*
 1022 *Cell* 7 (2005) 77-85.

1023 [30] C.L. Dalgard, H. Lu, A. Mohyeldin, A. Verma, Endogenous 2-oxoacids
 1024 differentially regulate expression of oxygen sensors, *Biochem J* 380 (2004) 419-424.

1025 [31] G.L. Wang, B.H. Jiang, E.A. Rue, G.L. Semenza, Hypoxia-inducible factor 1
 1026 is a basic-helix-loop-helix-PAS heterodimer regulated by cellular O₂ tension, *Proc*
 1027 *Natl Acad Sci U S A* 92 (1995) 5510-5514.

1028 [32] H.J. Knowles, R.R. Raval, A.L. Harris, P.J. Ratcliffe, Effect of ascorbate on
 1029 the activity of hypoxia-inducible factor in cancer cells, *Cancer Res* 63 (2003) 1764-
 1030 1768.

1031 [33] R.K. Bruick, S.L. McKnight, A conserved family of prolyl-4-hydroxylases that
 1032 modify HIF, *Science* 294 (2001) 1337-1340.

1033 [34] M. Ivan, K. Kondo, H. Yang, W. Kim, J. Valiando, M. Ohh, A. Salic, J.M.
 1034 Asara, W.S. Lane, W.G. Kaelin, Jr., HIFalpha targeted for VHL-mediated destruction
 1035 by proline hydroxylation: implications for O₂ sensing, *Science* 292 (2001) 464-468.

1036 [35] P. Jaakkola, D.R. Mole, Y.M. Tian, M.I. Wilson, J. Gielbert, S.J. Gaskell, A.
 1037 Kriegsheim, H.F. Hebestreit, M. Mukherji, C.J. Schofield, P.H. Maxwell, C.W. Pugh,
 1038 P.J. Ratcliffe, Targeting of HIF-alpha to the von Hippel-Lindau ubiquitylation complex
 1039 by O₂-regulated prolyl hydroxylation, *Science* 292 (2001) 468-472.

1040 [36] W.C. Hon, M.I. Wilson, K. Harlos, T.D. Claridge, C.J. Schofield, C.W. Pugh,
 1041 P.H. Maxwell, P.J. Ratcliffe, D.I. Stuart, E.Y. Jones, Structural basis for the
 1042 recognition of hydroxyproline in HIF-1 alpha by pVHL, *Nature* 417 (2002) 975-978.

1043 [37] N. Masson, C. Willam, P.H. Maxwell, C.W. Pugh, P.J. Ratcliffe, Independent
 1044 function of two destruction domains in hypoxia-inducible factor-alpha chains
 1045 activated by prolyl hydroxylation, *EMBO J* 20 (2001) 5197-5206.

1046 [38] J.H. Min, H. Yang, M. Ivan, F. Gertler, W.G. Kaelin, Jr., N.P. Pavletich,
 1047 Structure of an HIF-1alpha -pVHL complex: hydroxyproline recognition in signaling,
 1048 *Science* 296 (2002) 1886-1889.

1049 [39] V. Srinivas, L.P. Zhang, X.H. Zhu, J. Caro, Characterization of an
 1050 oxygen/redox-dependent degradation domain of hypoxia-inducible factor alpha (HIF-
 1051 alpha) proteins, *Biochem Biophys Res Commun* 260 (1999) 557-561.

1052 [40] K. Gradin, J. McGuire, R.H. Wenger, I. Kvietikova, M.L. Whitelaw, R. Toftgard,
 1053 L. Tora, M. Gassmann, L. Poellinger, Functional interference between hypoxia and
 1054 dioxin signal transduction pathways: competition for recruitment of the Arnt
 1055 transcription factor, *Mol Cell Biol* 16 (1996) 5221-5231.

1056 [41] P.J. Kallio, I. Pongratz, K. Gradin, J. McGuire, L. Poellinger, Activation of
 1057 hypoxia-inducible factor 1alpha: posttranscriptional regulation and conformational
 1058 change by recruitment of the Arnt transcription factor, *Proc Natl Acad Sci U S A* 94
 1059 (1997) 5667-5672.

1060 [42] S.M. Wood, J.M. Gleadle, C.W. Pugh, O. Hankinson, P.J. Ratcliffe, The role
 1061 of the aryl hydrocarbon receptor nuclear translocator (ARNT) in hypoxic induction of
 1062 gene expression. Studies in ARNT-deficient cells, *J Biol Chem* 271 (1996) 15117-
 1063 15123.

1064 [43] K.S. Hewitson, L.A. McNeill, M.V. Riordan, Y.M. Tian, A.N. Bullock, R.W.
 1065 Welford, J.M. Elkins, N.J. Oldham, S. Bhattacharya, J.M. Gleadle, P.J. Ratcliffe,
 1066 C.W. Pugh, C.J. Schofield, Hypoxia-inducible factor (HIF) asparagine hydroxylase is
 1067 identical to factor inhibiting HIF (FIH) and is related to the cupin structural family, *J*
 1068 *Biol Chem* 277 (2002) 26351-26355.

1069 [44] D. Lando, D.J. Peet, J.J. Gorman, D.A. Whelan, M.L. Whitelaw, R.K. Bruick,
 1070 FIH-1 is an asparaginyl hydroxylase enzyme that regulates the transcriptional
 1071 activity of hypoxia-inducible factor, *Genes Dev* 16 (2002) 1466-1471.

1072 [45] D. Lando, D.J. Peet, D.A. Whelan, J.J. Gorman, M.L. Whitelaw, Asparagine
 1073 hydroxylation of the HIF transactivation domain a hypoxic switch, *Science* 295
 1074 (2002) 858-861.

1075 [46] N. Sang, J. Fang, V. Srinivas, I. Leshchinsky, J. Caro, Carboxyl-terminal
 1076 transactivation activity of hypoxia-inducible factor 1 alpha is governed by a von
 1077 Hippel-Lindau protein-independent, hydroxylation-regulated association with
 1078 p300/CBP, *Mol Cell Biol* 22 (2002) 2984-2992.

1079 [47] C.C. Hudson, M. Liu, G.G. Chiang, D.M. Otterness, D.C. Loomis, F. Kaper,
 1080 A.J. Giaccia, R.T. Abraham, Regulation of hypoxia-inducible factor 1alpha
 1081 expression and function by the mammalian target of rapamycin, *Mol Cell Biol* 22
 1082 (2002) 7004-7014.

1083 [48] J.C. Anthony, F. Yoshizawa, T.G. Anthony, T.C. Vary, L.S. Jefferson, S.R.
 1084 Kimball, Leucine stimulates translation initiation in skeletal muscle of postabsorptive
 1085 rats via a rapamycin-sensitive pathway, *J Nutr* 130 (2000) 2413-2419.

1086 [49] G. D'Angelo, E. Duplan, P. Vigne, C. Frelin, Cyclosporin A prevents the
 1087 hypoxic adaptation by activating hypoxia-inducible factor-1alpha Pro-564
 1088 hydroxylation, *J Biol Chem* 278 (2003) 15406-15411.

1089 [50] R.C. Armstrong, Isolation and characterization of immature oligodendrocyte
 1090 lineage cells, *Methods* 16 (1998) 282-292.

1091 [51] C.J. Lynch, B. Halle, H. Fujii, T.C. Vary, R. Wallin, Z. Damuni, S.M. Hutson,
 1092 Potential role of leucine metabolism in the leucine-signaling pathway involving
 1093 mTOR, *Am J Physiol Endocrinol Metab* 285 (2003) E854-863.

1094 [52] R. Humar, F.N. Kiefer, H. Berns, T.J. Resink, E.J. Battegay, Hypoxia
 1095 enhances vascular cell proliferation and angiogenesis in vitro via rapamycin
 1096 (mTOR)-dependent signaling, FASEB J 16 (2002) 771-780.

1097 [53] E.J. Brown, M.W. Albers, T.B. Shin, K. Ichikawa, C.T. Keith, W.S. Lane, S.L.
 1098 Schreiber, A mammalian protein targeted by G1-arresting rapamycin-receptor
 1099 complex, Nature 369 (1994) 756-758.

1100 [54] D.M. Sabatini, H. Erdjument-Bromage, M. Lui, P. Tempst, S.H. Snyder,
 1101 RAFT1: a mammalian protein that binds to FKBP12 in a rapamycin-dependent
 1102 fashion and is homologous to yeast TORs, Cell 78 (1994) 35-43.

1103 [55] C.J. Sabers, M.M. Martin, G.J. Brunn, J.M. Williams, F.J. Dumont, G.
 1104 Wiederrecht, R.T. Abraham, Isolation of a protein target of the FKBP12-rapamycin
 1105 complex in mammalian cells, J Biol Chem 270 (1995) 815-822.

1106 [56] Z. Arany, L.E. Huang, R. Eckner, S. Bhattacharya, C. Jiang, M.A. Goldberg,
 1107 H.F. Bunn, D.M. Livingston, An essential role for p300/CBP in the cellular response
 1108 to hypoxia, Proc Natl Acad Sci U S A 93 (1996) 12969-12973.

1109 [57] J. Brismar, A. Aqeel, G. Brismar, R. Coates, G. Gascon, P. Ozand, Maple
 1110 syrup urine disease: findings on CT and MR scans of the brain in 10 infants, AJNR
 1111 Am J Neuroradiol 11 (1990) 1219-1228.

1112 [58] J.J. Riviello, Jr., I. Rezvani, A.M. DiGeorge, C.M. Foley, Cerebral edema
 1113 causing death in children with maple syrup urine disease, J Pediatr 119 (1991) 42-
 1114 45.

1115 [59] J. Josko, B. Gwozdz, H. Jedrzejowska-Szypulka, S. Hendryk, Vascular
1116 endothelial growth factor (VEGF) and its effect on angiogenesis, Med Sci Monit 6
1117 (2000) 1047-1052.
1118

Legends to Figures

Figure 1. BCAAs and BCKAs stabilize HIF-1 α in normoxia, reversible by ascorbate treatment.

A. Representative blots loaded with 20 μ g of whole-cell protein extracts from C6-ODD-GFP cells were analyzed for HIF-1 α , ODD-GFP, HIF-1 β , and phospho-S6R proteins (time=3hrs, n=4). **B, C, D.** Densitometric analyses of signals of pSer^{235/236}S6R, HIF-1 α/β , and ODD-GFP, respectively, are displayed in densitometry units normalized to non-treated controls. The specific conditions for treatments presented in the figure are hypoxia [1% O₂, 5% CO₂, and 94% N₂], KIC [2.0 mM], rapamycin [0.1 ng/mL], and ascorbate [100 μ M]. Statistical significances were determined by Tukey's HSD post hoc analysis (*p<0.05 compared to control; **p<0.001 compared to control; +p<0.05 compared to ascorbate treatment; ++p<0.001 compared to ascorbate treatment; ##p<0.001 compared to rapamycin treatment). **E, F.** C6-ODD-GFP cells were analyzed for VEGF protein by ELISA. Secreted protein was normalized to viable cell number determined by trypan blue uptake. The specific conditions for treatments presented in the figure are hypoxia [1% O₂, 5%CO₂, and 94% N₂], KIC [2.0 mM] (**E.**), leucine [2.0 mM] (**F.**), rapamycin [0.1 ng/mL], and ascorbate [100 μ M]. (time=12hrs, n=4). Statistical significances were determined by Tukey's HSD post hoc analysis (*p<0.05 compared to control; **p<0.001 compared to control; +p<0.05 compared to ascorbate treatment; ++p<0.001 compared to ascorbate treatment).

Figure 2. Dose-response for BCAAs and BCKAs on HIF-1 α level stabilization and functional roles in gene upregulation

A. Representative blots loaded with 20 μ g of whole-cell protein extracts from C6-ODD-GFP cells are shown for treatments corresponding to 0.5 increment increases on the log scale (time=3hrs, n=3). **B.** Densitometric analyses of HIF-1 α and phosphorylation status of Ser292-E1 α BCKDH. Data are normalized to non-treated control. **C.** Graph indicating non-treated control, positive control hypoxia [1% O₂, 5%CO₂, and 94% N₂], and dose-response of the BCKA KIC at 0.1, 0.5, 1.0, 2.5, and 5.0 mM (time=12hrs, n=4). **D.** Graph indicating non-treated control, hypoxia positive control, and dose-response of the BCAA leucine at 0.1, 0.5, 1.0, 2.5, and 5.0 mM (time=12hrs, n=4). Protein secretion levels shown in **C.** and **D.** are normalized to cell number. Statistical significances were determined by Dunnett's post hoc analysis (*p<0.05 compared to control; **p<0.001 compared to control).


Figure 3. KIC induces HIF-1 α stabilization in primary astrocyte-enriched rat cultures, reversible by ascorbate treatment.

Representative blots loaded with 20 μ g of whole-cell protein extracts from primary astrocyte-enriched cell cultures were analyzed for HIF-1 α and HIF-1 β proteins for non-treated control, hypoxia [1% O₂, 5%CO₂, and 94% N₂], KIC [0.1 and 1.0 mM], and KIC treatments with ascorbate [0.1 mM] (time=3hrs, n=4). **B.** Densitometric analyses of HIF-1 α / β signals displayed in densitometry units normalized to non-treated controls showing statistical significances as determined by Tukey's HSD post hoc analysis (*p<0.05 compared to control; **p<0.001 compared to control; *p<0.05

compared to 0.1 mM KIC ascorbate treatment; [#]p<0.05 compared to 1.0 mM KIC with ascorbate treatment). **C.** Primary astrocyte-enriched cultures were analyzed for VEGF protein secretion by ELISA for non-treated control, hypoxia [1% O₂, 5%CO₂, and 94% N₂], KIC [0.1 and 1.0 mM], and KIC treatments with ascorbate [0.1 mM] (time=12hrs, n=4). Protein levels were normalized to cell number for each sample prior to statistical analyses. Statistical differences were determined by Tukey's HSD post hoc analysis (***p<0.0001 compared to control; ⁺⁺⁺p<0.0001 compared to 0.1mM KIC with ascorbate treatment; ^{###}p<0.0001 compared to 1.0mM KIC with ascorbate treatment).

1174 **Figures**

1175 **Figure 1. BCAAs and BCKAs stabilize HIF-1α in normoxia, reversible by**
1176 **ascorbate treatment**




1177

1178

1179
1180

Figure 2. Dose-response for BCAAs and BCKAs on HIF-1 α level stabilization and functional roles in gene upregulation




1181

1182

1183
1184

Figure 3. KIC induces HIF-1 α stabilization in primary astrocyte-enriched rat cultures, reversible by ascorbate treatment



1185

1186

1187

1188 **Alterations to branched-chain amino acid metabolism increase *in vitro***
1189 **malignant characteristics of C6 glioma cells**

1190 Jeremy HENRIQUES^{*1}, Thomas MCFATE¹, John TRENTINI¹, Emily HATHAWAY²,
1191 Michael SCHELL^{1,3}, Nam Ho JEOUNG⁴, Robert A HARRIS⁴, and Ajay VERMA^{1,5}

1192 ¹Neuroscience Program, Uniformed Services University of the Health Sciences,
1193 Bethesda, MD

1194 ²School of Medicine, Uniformed Services University of the Health Sciences,
1195 Bethesda, MD

1196 ³Department of Pharmacology, Uniformed Services University of the Health
1197 Sciences, Bethesda, MD

1198 ⁴Department of Biochemistry and Molecular Biology, The Indiana University School
1199 of Medicine, Indianapolis, ID

1200 ⁵Department of Neurology, Uniformed Services University of the Health Sciences,
1201 Bethesda, MD

1202 ^{*}To whom correspondence should be addressed

1203 Jeremy Henriques

1204 Neuroscience Program

1205 Uniformed Services University of the Health Sciences

1206 4301 Jones Bridge Road

1207 Bethesda, MD 20814

1208 Telephone: (301) 295-3840

1209 Fax: (301) 295-3825

1210 e-mail: henriques.jeremy@gmail.com

1211

1212 Abbreviations: BCAA, branched-chain amino acid; BCKA, branched-chain α -
1213 ketoacid; BCKDC, branched-chain α -ketoacid dehydrogenase complex; BCKDH,
1214 branched-chain α -ketoacid dehydrogenase; BDK, BCKDH kinase; BDP, BCKDH
1215 phosphatase; HIF, hypoxia inducible factor; KIC, 2-ketoisocaproate; KIV, 2-
1216 ketoisovalerate; KMV, 2-keto-3-methyl valerate; mTOR, mammalian target of
1217 rapamycin; PDH, pyruvate dehydrogenase; PDK, PDH kinase

1218

1219 **Summary**

1220 Branched-chain amino acids (BCAAs) participate in several biochemical pathways
1221 and are emerging as novel signal-initiating molecules. BCAAs are exclusively
1222 metabolized through the branched-chain α -ketoacid dehydrogenase complex
1223 (BCKDC), a mitochondrial macroenzyme complex, which is regulated through
1224 reversible phosphorylation. Cancer cells are in a state of hyperactivity utilizing
1225 metabolic substrates for energy (catabolism), for synthesis of new molecules
1226 (anabolism), and/or replenishing pools of intermediary metabolites (anaplerosis).
1227 Here we sought to better understand the impact of BCAA metabolism on cancer
1228 cells by altering the activity state of the BCKDC. We successfully created a shRNA
1229 protein knockdown system for the BCKDC kinase (BDK) in C6 glioblastoma cells,
1230 thus preventing phosphorylation and increasing activity of BCKDC. We found BDK
1231 knockdown cells had greater proliferation, invasion, and migration rates compared
1232 with control vector and wild type cell lines. We also found an increased lactate
1233 output for the BDK knockdown cells and pyruvate dehydrogenase complex (PDC)

1234 phosphorylation, both are hallmarks of “The Warburg Effect”. These data indicate
1235 compensatory changes occur to retain metabolic homeostasis as cancer cells are
1236 forced to promote BCAA oxidation. The result is increased progressive *in vitro* cell
1237 behavior. Our conclusion is that these two metabolic systems, BCKDC and PDC,
1238 may not act independently of one another in cancer cells.

1239 *Introduction*

1240 Branched-chain amino acids (BCAAs: leucine, isoleucine, and valine)
1241 comprise a subgroup of essential amino acids that cannot be synthesized *de novo*
1242 and have multipotent roles in both biochemistry and signaling biology. Leucine is
1243 particularly potent, if not singularly effective, in many signaling pathways (eg,
1244 translational activation (Mordier and others 2000), insulin synthesis and secretion
1245 (Lambert and others 1986), inhibition of autophagy (Mordier and others 2000), and
1246 hypothalamic satiety signaling (Cota and others 2006)). BCAAs can also be
1247 deaminated to α -keto acids and metabolized for fatty acid synthesis and β -oxidation
1248 (Greenberg and Reaven 1966; Noda and Ichihara 1974; Noda and Ichihara 1976).
1249 Owing to these many biological pathways open to BCAAs, it is peculiar that their
1250 irreversible metabolism is known to be governed by one multi-enzyme complex, the
1251 branched-chain α -keto acid dehydrogenase complex (BCKDC).

1252 The BCKDC is composed of E1, E2 and E3 subunits, similar to other
1253 dehydrogenases, pyruvate dehydrogenase complex (PDC) and α -ketoglutarate
1254 dehydrogenase complex (Harris and others 1995). First step metabolism proceeds
1255 through the heterodimeric E1 subunit comprised of two α subunits and two β
1256 subunits (Danner and others 1978; Parker and Randle 1978; Pettit and others 1978;
1257 Roberts and Sokatch 1978). The BCKDC is active when the serine residue 292 on
1258 the E1 α subunit is dephosphorylated and is inactive upon phosphorylation. This
1259 reversible phosphorylation is regulated by a specific kinase (BDK) (Popov and
1260 others 1992; Shimomura and others 1990) and phosphatase (BDP) (Damuni and
1261 others 1984; Damuni and Reed 1987). The biochemical processes occurring through

the BCKDC are irreversible. Therefore, the careful regulation of the BCKDC is the most important step in BCAA metabolism.

In normal tissues, BDK and BDP carefully control the activity of the BCDKC. This control regulates the amount of free BCAAs available not only for catabolic energy production, but also anabolic and signaling requirements. Anabolic demands for BCAAs include fatty acid synthesis and single amino acids used in the synthesis of proteins. In the absence of sufficient amounts of BCAAs, the translation is unable to occur. This is not only owing to the lack of BCAAs, but also the signaling roles BCAAs have in initiating translation. Leucine in particular has been shown to initiate translational machinery indirectly by stimulating insulin secretion and directly via the mammalian target of rapamycin (mTOR) (Kimball and others 1999). The fate of BCAAs amidst these different pathways is carefully controlled in the normal state. However, it has been shown that when tissues and cells are in a state of hypermetabolism (ie, cancer (Baracos 2000; Baracos and Mackenzie 2006)), the free pool of BCAAs is used up rapidly for energy and protein synthesis requirements. Increasing energy demands of tumor-bearing rats have been shown to increase leucine oxidation measured by radio-labeled CO₂ release (Costelli and others 1995), indicating a systemic depletion of leucine in the host organism through BCKDC activity. As a compensatory mechanism to refill the rapidly depleting free BCAA pool in patients with severe, invasive cancer, skeletal muscle, which is rich in BCAAs, is degraded.

The compensatory increases in free BCAAs, high protein turnover rates, and high oxidation rates cause increased BCKA levels, at least transiently. Our lab has

recently shown high levels of BCKAs have signaling properties through the hypoxia inducible factor (HIF) (manuscript in submission). This survival factor is known to play an important role in cancer, particularly in altering transcription of essential proteins for proliferation, invasion, and compensatory metabolism (reviewed in (Ke and Costa 2006; Semenza 2007)). HIF specifically alters glucose metabolism by increasing glucose transporters, PDC kinase (PDK), and lactate dehydrogenase (LDH), to name just a few. These actions drive anaerobic glycolysis which is advantageous since it has been shown that tumors exist in a hypoxic environment. However, this effect has been shown in the presence of oxygen, an observation known as the Warburg Effect (Warburg 1930; Warburg 1956). Regardless of the many changes that occur as a result of HIF activation in cancer, our lab has recently shown that PDK activity is directly responsible for increasing cancerous phenotypes (McFate and others 2008). PDK acts to shut down PDC activity the same way BDK acts to shut down BCKDC. Given that PDK activity can be manipulated to alter cancer phenotypes, we hypothesized that altering BDK activity would also alter cancer phenotypes.

In this publication, we show that utilizing stably transfected shRNA knockdown system of BDK in cancer cells, the phenotypes of the cells are altered. However, these data also suggest that by altering the BCKDC system, metabolic compensation occurs to the point where PDC activity is affected. These data indicate that the dehydrogenases PDC and BCKDC may not act independently of one another.

1307 **Experimental Methods**

1308 All chemicals were purchased from Sigma-Aldrich unless otherwise stated. Cell
1309 culture products were purchased from GIBCO.

1310 **Cell culture and Chemical Treatments**

1311 Cells were cultured in Dulbecco's Modified Eagle Media (DMEM, Invitrogen)
1312 supplemented with 10% fetal bovine serum, 1% penicillin-streptomycin, and 1.5
1313 mg/mL G418 (Gibco). G418 was used to select transgenically altered cells. Cells
1314 were cultured under 37°C incubation and in 21% O₂, 5%CO₂, and 74% N₂. Chemical
1315 treatments were performed in Kreb's Saline Buffer and incubation times are as
1316 indicated. Each chemical treatment was performed by dilution from 100X stock in
1317 Kreb's saline buffer. For cell hypoxia treatment, the culture dishes were sealed in a
1318 modular incubator chamber, flushed with gas containing 1% O₂, 5%CO₂, and 94%
1319 N₂ for 5 min, and incubated in this environment at 37°C for the indicated times.

1320 **Generation of Transgenic C6 Glioma Cell Lines for BDK Knockdown**

1321 An shRNA knockdown vector for transfection into C6 glioma cells specific to BDK
1322 was obtained from SuperArray. The cells with shRNA BDK are referred to as shK
1323 (shRNA BCKD-Kinase) from hereafter. The SuperArray kit provided a standard
1324 control vector consisting of a scrambled DNA sequence cloned into the identical
1325 vector for quality and specificity of effects that we also used and is referred to as
1326 CVC6 (control vector for C6 cells) hereafter. CVC6 and shK cells were treated
1327 identically through the transfection procedures. The cells were transfected as
1328 described in the product manual. Briefly, vectors were grown in *E.coli* JM109
1329 (Stratagene). DNA was purified by a commercial miniprep kit (Qiagen). DNA was

then transfected into C6 cells with Lipofectamine 2000 reagent (Invitrogen) and incubated in fresh DMEM with 10% FBS following transfection. After 48 hrs recovery time, transfected cells were selected by adding 1.5 mg/mL G418 (Gibco) in complete DMEM. Media was changed as cells died and colonies were allowed to grow until large enough to pluck and grow independently. Once cells were grown and verified through Western blotting of BDK, they were used as stable cell stocks to ensure clonal cell lines. shK and CVC6 cell lines were incubated as previously described for wild type in the “**Cell Culture and Chemical Treatments**” section.

Cell counting

Cell were washed and trypsinized for cell counting procedures for ELISA normalization. Trypsin was inactivated by adding equal amounts of media with 10% FBS. Five hundred μ L of suspensions of each preparation were then placed in a Vi-Cell Trypan Blue automated counter for cell counting.

Western Blot and Densitometry Analyses

Cells were washed three times with cold PBS. Appropriate amount of lysis buffer containing RIPA buffer (Bioworld), 1% SDS, and 1X protease inhibitor cocktail (Roche) then scraped. Cell remnants were then collected in 1.5 ml microcentrifuge tubes. The cell material was sonicated for 25 seconds at 50 Hz, then placed in rack at room temperature. Cell lysates were then spun for 5 minutes at 12,000 g and the supernatants were transferred to fresh tubes. Protein levels were determined by the BCA method of analysis (Pierce). Proteins were separated on 4-12% Bis-Tris SDS-polyacrylamide gradient gels (Invitrogen) and transferred to nitrocellulose membranes (Invitrogen). Membranes were blocked using 5% goat or horse serum

(Vecotr Labs) in Tris buffered saline with 0.1% Tween-20. Antibodies used were anti-phospho-Ser292-E1 α BCKDH (generous gift from C. Lynch, 1:20,000), anti-E1 α BCKDH (generous gift from C. Lynch, 1:1000), anti-BDK 1:2000 (generous gift from Dr. R. Harris), E1 β /E2 antisera 1:2000 (generous gift from Dr. R. Harris), anti-phospho-Ser293-E1 α PDC 1:5000 (Novus), anti-E1 α PDC 1:2000 (Novus) and anti- β actin 1:10,000 (AbCam). Protein bands were visualized by enhanced chemiluminescence (Pierce) using either Kodak film and developer or CCD luminescence camera (Fuji Film). Signals were quantified using densitometry in ImageJ (Wayne Rasband, NIH). Phosphoproteins were normalized to total protein amounts unless otherwise stated. Graphs indicate arbitrary units on the y-axes derived from signals normalized to non-treated or wild type controls by direct comparative ratios. These normalizations were necessary to control for protein loading.

Proliferation assays

One hundred thousand cells were plated in 6-well culture dishes to a volume of 2 mL of DMEM media solution with 10% FBS and 1% penicillin-streptomycin. At time points, cells were counted as described in the “*Cell Counting*” procedure. The data are presented as either raw cell numbers or relative proliferation rates normalized to wild type controls.

Migration and Invasion Assays

Migration experiments were performed using BD Falcon cell culture inserts with 8 micron pores (BD Biosciences). Cell invasion experiments were performed using 24-well Biocoat Matrigel™ invasion chambers with an 8- μ m pore polycarbonate filter

according to the manufacturer's instructions (BD Biosciences) Fifty thousand cells were plated in migration and invasion inserts to a volume of 0.5 mL of DMEM solution. The inserts were then placed in 24-well culture dishes containing 0.75 mL DMEM solution with 1% FBS and 1% penicillin-streptomycin. Cells were incubated at 37 degrees Celsius, 1% O₂, 5.0% CO₂ and 94% N₂ for 72 hr. Cells that remained inside the insert after 48 hours were thoroughly wiped with a cotton swab and invading cells were fixed and stained using Diff-Quick Stain Solution (Dade Behring). Images were then taken with a Canon CoolSNAP digital camera in Metamorph imaging program under light microscopy. Cell number was determined by marking individual cells with spots in the nucleus in ImageJ imaging software and using the quantifier for spot counting. Five images were taken, counted and averaged from each well insert to ensure adequate sampling.

Colony Formation Assay

Ten thousand cells from each cell line were placed in 1 ml of DMEM with 0.3% low-melting agarose (soft agar) and 10% FBS, and overlaid onto 1 ml/well of DMEM with 0.5% agarose and 10% FBS. Cells were then incubated for 18 days at 37 C in 5.0% CO₂. At the endpoint, wells were placed under light microscope with COOLSnap Kodak camera. Five images were taken, counted and averaged from each well insert to ensure adequate sampling. Colonies were counted by identification with exclusion parameters at ~10 nm. Colony diameters were determined by importing tiff images in ImageJ software. A line was then drawn from end to end of colony going through the center and ImageJ quantified diameter.

CMA Analysis of Lactate

Cells cultured and samples of media were taken at indicated times and frozen at -80°C. Cell numbers were then obtained as described previously in the "*Cell Counting*" section. Media samples were thawed, vortexed and centrifuged to eliminate air bubbles. 50 µL of media samples were used to measure metabolite levels. Lactate was measured using a CMA 600/microdialysis analyzer (CMA Microdialysis AB). Resulting data were normalized to cell number for each sample.

BCKDC Activity Assay by Radiolabeled CO₂ Capture

The following analysis was performed by collaborators (Jeoung and others 2006). Briefly, cells were plated in culture flasks (1 x 10⁶ cell/flask for wild type cells, 0.75 x 10⁶ cell/flask for CVC6 and shk cells because growth rates were different.) Cells were cultured until growth reached 90% confluency. Cells were then washed twice with PBS (room temperature). Five mL of Krebs-Henseleit buffer containing 5 mM glucose, 1 mM ¹⁴C-valine (specific activity; 150 uCi/mmol), 0.2 % BSA, and 1 mU/mL insulin. The air was flushed with 95% O₂ and 5% N₂, then the bottle was closed with a rubber stopper. Experimental incubation time was 1 hr at 37°C. Three-hundred µL of 60% perchloric acid was added to stop the reaction process. ¹⁴CO₂ was collected into the center well. In order to check for KIV production from valine, 1 mL of stopped solution was taken out and put into a scintillation vial. The rubber stopper was then replaced and 350 µL of 30% H₂O₂ was added and incubated for 30 min. Radioactivity in the center well was then counted. Data are expressed as nmol of oxidized valine determined to be in equimolar proportions with captured ¹⁴CO₂ and normalized to cell number.

ATP bioluminescence assay

Cells were grown under culture conditions. Cells were detached from the culture plates with trypsin and suspended in media for counting. Media volumes were adjusted to a range of 10^5 to 10^8 cells/mL. Boiling 100 mM Tris, 4 mM EDTA, pH 7.75 was added (9:1 ratio to sample volume) and incubated for 2 min at 100°C. Samples were then centrifuged at 1000g for 60 sec. The supernatants were transferred to clean tubes. Luciferase reagent was then added to the samples and bioluminescence was recorded after 10, 15, and 20 seconds. The highest values were taken and normalized to viable cell number established from left over cell suspensions for each sample.

Statistical Analyses

Statistical analyses ANOVA, Tukey's HSD and Fisher's PLSD *post hoc* tests were performed using STATview statistical software. Where it was necessary to compare treatment only to control groups, ANOVA and Dunnett's *post hoc* analyses were performed in SPSS statistical software.

Results

A specific BDK knockdown using stably transfected shRNA in C6 glioma cells proves viable in reducing BCKDC phosphorylation and activity

C6 glioma cells have a high basal level of BCKDC phosphorylation. It is unclear what role the metabolism of BCAAs and BCKAs has in cancer biochemistry. Previous reports are perfunctory and contradictory occurring mostly in tumor-bearing animals, but also in humans, leaving the question open to further and more specific investigation. We chose to manipulate an *in vitro* system of cancer to identify any key differences observed when the BCKDC activity state was constitutively active.

These C6 cells were used to generate stable-shRNA BDK knockdown cells (shK) and stable-control vector (CVC6) cell lines. The shK and CVC6 cells generated were analyzed via Western blot for BDK and phosphorylated Ser292-BCKDC E1 α to verify that there had reduced and comparable signals, respectively, as compared to C6 wild type (Fig. 1A). Densitometry of Fig. 1A Western blots are also shown in Fig. 1B. The shK cell line had a decreased BDK signal compared with both wild type ($p<0.0001$) and CVC6 ($p<0.0001$). There were no differences between wild type and CVC6 cell lines in BDK signal. Phosphorylated Ser292-BCKDC E1 α levels for shK cells were non-detectable, whereas wild type and CVC6 cell lines had comparable signals. For normalization of phosphorylated signals of proteins, the phosphorylated signal is divided by the total protein signal and expressed as a ratio. The shK cells had a decreased ratio of phosphorylated Ser292-BCKDC E1 α from wild type ($p<0.0001$) and CVC6 ($p<0.0001$), whereas no differences were detected between wild type and CVC6 cells. The activity of the E1 α -BCKDC was generously performed by Nam Ho Jeung, a collaborator, and shows shK cells have an increased activity of the E1 α subunit as compared to wild type and CVC6 ($p<0.05$ for both comparisons; Fig. 1B).

E1 β and E2 Subunits of BCKDC in the Different Cell Lines

BCKDC subunits E1 β and E2 were analyzed by Western blot and densitometry (Fig. 2A and B, respectively) to see if any disruption to normal levels of these proteins resulted from the transgenic alterations to BDK translation. The E1 β subunit is the heterodimeric partner to the E1 α that composes the E1 subunit. There were no statistical differences of E1 β signal between shK cells when compared to

either the wild type or CVC6. No differences were observed between wild type and CVC6. However, the signals from the blot of the E2 subunits of each cell line showed increased signals for shK cells compared with wild type ($p<0.05$). No differences were present between either the shK and CVC6 cell lines, or the wild type and CVC6 cell lines.

BDK Knockdown Cell Line Shows Increased Proliferation from Controls

Cell proliferation was compared for the cell lines over 72 hrs, the data from which are depicted in Fig. 3A. The number of viable shK cells was significantly increased compared with wild type for 24 ($p<0.001$), 48 ($p<0.0001$) and 72 hrs ($p<0.0001$). The number of viable shK cells was also significantly increased compared with CVC6 for 24 ($p<0.001$), 48 ($p<0.0001$) and 72 hrs ($p<0.0001$). The average proliferation rate of shK cells across 24, 48 and 72 hr time points was 137.63% of wild type and 136.82% of CVC6 cells. There were no differences between wild type and CVC6 cells for any time points. To investigate if the increased proliferation was due to growth factor signaling, we replicated the proliferation assay in serum-free media (Fig. 3B). The results show increased shK cell proliferation with respect to wild type at 48 ($p<0.05$) and 72 hrs ($p<0.05$), but not at 24 hrs. The number of shK cells was also significantly increased from CVC6 at 48 ($p<0.01$) and 72 hrs ($p<0.05$), but not 24 hrs. The average proliferation rate of shK cells across 24, 48, and 72 hr time points was 119.65% of wild type and 122.08% of CVC6 cells. There were no differences between wild type and CVC6 cells for any time points.

Metastatic Behaviors Increase in BDK Knockdown Cells Relative to Controls

To investigate if the increased rates of proliferation observed in the BDK knockdown cells were indicative of other malignant phenotypes, we measured the migration of cells through perforated culture inserts. This tests the mobility of cancer cells commonly seen in *in vivo* systems. We observed increased migratory rates for shK cells as compared to wild type ($p<0.0001$) and CVC6 ($p<0.0001$) cells, as shown in Fig. 4A and quantified in 4D.

Cell invasiveness, another metastatic phenotype commonly found in malignant cancerous tissue, was also analyzed. Invasion assays differed from migration only in that the cells were cultures in perforated culture inserts filled with extracellular matrix matrigel to mimic the extracellular environment. The results show increased invasiveness for shK cells as compared to wild type ($p<0.0001$) and CVC6 ($p<0.0001$) cells, as shown in Fig. 4B and quantified in 4E.

Cell lines were mixed with a soft agar DMEM solution and plated in 6 well dishes. Incubation of cells consisted of 3 weeks under culture conditions. Upon termination of the experiment, images were taken of cell colonies and analyzed as described in the *Experimental Methods* section. There were no differences in colony numbers between the three cell lines (data not shown). However, the size of the shK cell colonies was significantly increased with respect to wild type ($p<0.0001$) and CVC6 ($p<0.0001$) colonies, as shown in Fig. 4C and quantified in 4F.

Phosphorylated Characteristics of PDC, a Family member of the Heterodimeric Multimer Complex also Implicated in Metastatic Cancers

We next wondered if other mitochondrial dehydrogenases were affected by the BDK knockdown. Biochemical pathways in metabolism are tightly regulated

according to energy demands of the cell; alterations to one pathway may affect the another. Our lab had previously investigated PDC activity and its relation to cancerous phenotypes, classically termed “the Warburg Effect” (McFate and others 2008). The more aggressive cancerous phenotypes presented here were unexpected, leading us to investigate PDC metabolic activity. Cells were harvested and analyzed through Western blotting for E1 α PDC phosphorylation state. The phosphorylation state of Ser 293-E1 α PDC is shown with the total PDC E1 α signal beneath as loading controls (Fig. 5A). Densitometry of Western blots in Fig. 5A is represented in Fig. 5B. The PDC E1 α phosphorylation ratios are increased for the shK cell line when compared to both wild type ($p<0.05$) and CVC6 ($p<0.05$) cell lines.

Lactate Production as an Indicator of Warburg Characteristics

If the Warburg hypothesis is correct, then the BDK knockdown cells, which exhibit a more aggressive metastatic phenotype, should release more lactate into the culture medium. To test this, media samples were taken at 6, 12, 24, and 48 hrs for lactate analysis using CMA. Cell number was then determined as previously described. CMA analysis of the cultured media showed differences in lactate produced for the cell lines (Fig. 5C). The shK cells had increased lactate production with respect to wild type cells at 12 ($p<0.01$), 24 ($p<0.05$) and 48 hours ($p<0.05$). The shK cells also had increased lactate production with respect to CVC6 cells at 12 ($p<0.01$), 24 ($p=0.01$) and 48 hours ($p<0.05$).

Discussion

BCAA metabolism has been underappreciated in cancer biochemistry. Here we propose that shunting BCAA metabolism through the BCKDC affects overall cellular biochemistry in cancer cells. Using stably transfected shRNA, we successfully knocked down the translation of the BDK (shK cells), the kinase responsible for inactivation of the BCKDC, thus leaving the complex in a constitutively active state. Both phenotypical and biochemical changes were observed in the transgenically altered cancer cells. We found that shK cells were more proliferative, migratory, invasive, and able to form colonies in soft agar, an *in vitro* technique which shows cellular ability to form cancerous nodules. We also found that increased BCKDC activity in shK cells resulted in decreased activation of the PDC, another key mitochondrial metabolic multi-enzyme complex, and consequential increased lactate secretion from these cells. These data suggest that enhancing BCAA and BCKA metabolism through the BCKDC acts to modify cellular biochemistry so that PDC activity is decreased, which results in increased lactate production and increased invasive cancer phenotypes, historically dubbed the “Warburg Effect”.

Cancerous cells are in a constant state of hypermetabolism and accelerated growth. As such, their metabolic needs greatly differ from normal cells. Changes to their intracellular biochemistry cause appreciable differences to their behaviors and extracellular signaling. When cellular processes become dysregulated, like in cancer, the cell then has an altered transcriptional milieu from its genes in response.

1556 These changes are largely indicative of evolutionarily preserved survival and
1557 adaptation mechanisms.

1558 Recently, our lab has shown that increased levels of BCAAs and BCKAs can
1559 stimulate signaling mechanisms involving HIF, an evolutionarily conserved
1560 transcription factor which is normally activated by hypoxic stress conditions
1561 (manuscript in submission). Activation of the HIF pathway has been shown in cancer
1562 cells, where the level of increase above baseline corresponded to the degree to
1563 which invasive and metastatic characteristics were increased. These increased
1564 cancerous phenotypes have been shown to involve specific factors including VEGF,
1565 GLUT1, etc (reviewed in (Ke and Costa 2006)). However, increased activation of
1566 HIF has also been shown to be directly stimulated by increased levels of glycolytic
1567 metabolites, particularly pyruvate, OAA, succinate, and fumarate (Isaacs and others
1568 2005; Lu and others 2005; Selak and others 2005). HIF activation has been shown
1569 to increase glycolysis and decrease mitochondrial respiration, primarily through
1570 increased PDK expression, which leads to decreased PDC activity, a buildup of
1571 pyruvate, and, consequently, an increase in lactate production (Koukourakis and
1572 others 2005).

1573 Our lab has recently published data increasing the activity of PDC, the
1574 metabolic regulator of aerobic metabolism at the crux between glycolysis and
1575 cellular respiration, by PDK knockdown. These data showed increasing PDC activity
1576 through knockdown of PDK, HIF levels decreased, as did cancerous phenotypes of
1577 invasion, migration, and tumor size (McFate and others 2008).

1578 Our data show a decrease in presence and activity of BDK results in
1579 increased malignancy. These findings are contrary to our original hypothesis.
1580 However, perhaps a more comprehensive understanding has been reached in that
1581 the regulation of just one metabolic pathway is not always sufficient, as was shown
1582 by McFate et al (2008) and others (McFate and others 2008). Biochemistry of
1583 cancerous, as well as normal, cells requires a coordination of metabolic complexes.
1584 Our intent was to investigate whether altered BCKDC activity was capable of
1585 singularly altering the behavior of cancerous cells and cancer biochemistry in a way
1586 that would make it a new target for cancer therapy research, similar to PDC.
1587 However, it seems that there is no direct translation of metabolic complex to
1588 therapeutic target.

1589 In summary, BCAAs are metabolized through one multienzyme complex,
1590 BCKDC. In this report, we used an shRNA knockdown approach of BDK to cause
1591 hyperactivity of BCKDC. Our hypothesis was that by increasing BCKDC activity, the
1592 free amino acid pools would be depleted in cancer cells, which would limit their
1593 growth rates. The data show a successful knockdown of BDK to result in increased
1594 activation of BCKDC. However, the behavior of these cells increased in cancerous
1595 characteristics, such as proliferation, migration, invasion, and colony size. We further
1596 investigated whether these results were due to a coordination of metabolic
1597 complexes, namely BCDKC and PDC. These data show that alterations to BCKDC
1598 activity does indeed have an effect on PDC activity. The nature of this effect is one
1599 that drives PDC to become less active, a characteristic shown by our lab and others
1600 as causing increased cancerous behaviors. It seems that BCAA control is not solely

responsible for cancer behaviors, but these unexpected findings do, however,
warrant a closer look at BCAAs in cancer biochemistry.

Acknowledgements: This work was funded by NIH grants NS73814 and CA113506
to AV.

References

- S. Mordier, C. Deval, D. Bechet, A. Tassa, and M. Ferrara, Leucine limitation
induces autophagy and activation of lysosome-dependent proteolysis in
C2C12 myotubes through a mammalian target of rapamycin-independent
signaling pathway. J Biol Chem 275 (2000) 29900-6.
- D.G. Lambert, K. Hughes, and T.W. Atkins, Insulin release from a cloned hamster B-
cell line (HIT-T15). The effects of glucose, amino acids, sulphonylureas and
colchicine. Biochem Biophys Res Commun 140 (1986) 616-25.
- D. Cota, K. Proulx, K.A. Smith, S.C. Kozma, G. Thomas, S.C. Woods, and R.J.
Seeley, Hypothalamic mTOR signaling regulates food intake. Science 312
(2006) 927-30.
- R. Greenberg, and G. Reaven, The effect of L-leucine on hepatic glucose formation.
Pediatrics 37 (1966) 934-41.

1620 C. Noda, and A. Ichihara, Control of ketogenesis from amino acids. II. Ketone bodies
 1621 formation from alpha-ketoisocaproate, the keto-analogue of leucine, by rat
 1622 liver mitochondria. J Biochem 76 (1974) 1123-30.

1623 C. Noda, and A. Ichihara, Control of ketogenesis from amino acids. IV. Tissue
 1624 specificity in oxidation of leucine, tyrosine, and lysine. J Biochem 80 (1976)
 1625 1159-64.

1626 R.A. Harris, K.M. Popov, Y. Zhao, N.Y. Kedishvili, Y. Shimomura, and D.W. Crabb,
 1627 A new family of protein kinases--the mitochondrial protein kinases. Adv
 1628 Enzyme Regul 35 (1995) 147-62.

1629 D.J. Danner, S.K. Lemmon, and L.J. Elsas, 2nd, Substrate specificity and
 1630 stabilization by thiamine pyrophosphate of rat liver branched chain alpha-
 1631 ketoacid dehydrogenase. Biochem Med 19 (1978) 27-38.

1632 P.J. Parker, and P.J. Randle, Partial purification and properties of branched-chain 2-
 1633 oxo acid dehydrogenase of ox liver. Biochem J 171 (1978) 751-7.

1634 F.H. Pettit, S.J. Yeaman, and L.J. Reed, Purification and characterization of
 1635 branched chain alpha-keto acid dehydrogenase complex of bovine kidney.
 1636 Proc Natl Acad Sci U S A 75 (1978) 4881-5.

1637 C.M. Roberts, and J.R. Sokatch, Branched chain amino acids as activators of
 1638 branched chain ketoacid dehydrogenase. Biochem Biophys Res Commun 82
 1639 (1978) 828-33.

1640 K.M. Popov, Y. Zhao, Y. Shimomura, M.J. Kuntz, and R.A. Harris, Branched-chain
 1641 alpha-ketoacid dehydrogenase kinase. Molecular cloning, expression, and
 1642 sequence similarity with histidine protein kinases. J Biol Chem 267 (1992)
 1643 13127-30.

1644 Y. Shimomura, N. Nanaumi, M. Suzuki, K.M. Popov, and R.A. Harris, Purification
 1645 and partial characterization of branched-chain alpha-ketoacid dehydrogenase
 1646 kinase from rat liver and rat heart. Arch Biochem Biophys 283 (1990) 293-9.

1647 Z. Damuni, M.L. Merryfield, J.S. Humphreys, and L.J. Reed, Purification and
 1648 properties of branched-chain alpha-keto acid dehydrogenase phosphatase
 1649 from bovine kidney. Proc Natl Acad Sci U S A 81 (1984) 4335-8.

1650 Z. Damuni, and L.J. Reed, Purification and properties of the catalytic subunit of the
 1651 branched-chain alpha-keto acid dehydrogenase phosphatase from bovine
 1652 kidney mitochondria. J Biol Chem 262 (1987) 5129-32.

1653 S.R. Kimball, L.M. Shantz, R.L. Horetsky, and L.S. Jefferson, Leucine regulates
 1654 translation of specific mRNAs in L6 myoblasts through mTOR-mediated
 1655 changes in availability of eIF4E and phosphorylation of ribosomal protein S6.
 1656 J Biol Chem 274 (1999) 11647-52.

1657 V.E. Baracos, Regulation of skeletal-muscle-protein turnover in cancer-associated
 1658 cachexia. Nutrition 16 (2000) 1015-8.

1659 V.E. Baracos, and M.L. Mackenzie, Investigations of branched-chain amino acids
 1660 and their metabolites in animal models of cancer. J Nutr 136 (2006) 237S-
 1661 42S.

1662 P. Costelli, M. Llovera, C. Garcia-Martinez, N. Carbo, F.J. Lopez-Soriano, and J.M.
 1663 Argiles, Enhanced leucine oxidation in rats bearing an ascites hepatoma
 1664 (Yoshida AH-130) and its reversal by clenbuterol. Cancer Lett 91 (1995) 73-8.

1665 Q. Ke, and M. Costa, Hypoxia-inducible factor-1 (HIF-1). Mol Pharmacol 70 (2006)
 1666 1469-80.

1667 G.L. Semenza, HIF-1 mediates the Warburg Effect in clear cell renal carcinoma. J
 1668 Bioenerg Biomembr 39 (2007) 231-4.

1669 O. Warburg, The metabolism of tumors. Constable and Company, Ltd. (1930).

1670 O. Warburg, On the origin of cancer cells. Science 123 (1956) 309-14.

1671 T. McFate, A. Mohyeldin, H. Lu, J. Thakar, J. Henriques, N.D. Halim, H. Wu, M.J.
 1672 Schell, T.M. Tsang, O. Teahan, S. Zhou, J.A. Califano, N.H. Jeoung, R.A.
 1673 Harris, and A. Verma, Pyruvate dehydrogenase complex activity controls
 1674 metabolic and malignant phenotype in cancer cells. J Biol Chem 283 (2008)
 1675 22700-8.

1676 N.H. Jeoung, P. Wu, M.A. Joshi, J. Jaskiewicz, C.B. Bock, A.A. Depaoli-Roach, and
 1677 R.A. Harris, Role of pyruvate dehydrogenase kinase isoenzyme 4 (PDHK4) in
 1678 glucose homeostasis during starvation. Biochem J 397 (2006) 417-25.

1679 J.S. Isaacs, Y.J. Jung, D.R. Mole, S. Lee, C. Torres-Cabala, Y.L. Chung, M. Merino,
 1680 J. Trepel, B. Zbar, J. Toro, P.J. Ratcliffe, W.M. Linehan, and L. Neckers, HIF
 1681 overexpression correlates with biallelic loss of fumarate hydratase in renal
 1682 cancer: novel role of fumarate in regulation of HIF stability. *Cancer Cell* 8
 1683 (2005) 143-53.

1684 H. Lu, C.L. Dalgard, A. Mohyeldin, T. McFate, A.S. Tait, and A. Verma, Reversible
 1685 inactivation of HIF-1 prolyl hydroxylases allows cell metabolism to control
 1686 basal HIF-1. *J Biol Chem* 280 (2005) 41928-39.

1687 M.A. Selak, S.M. Armour, E.D. MacKenzie, H. Boulahbel, D.G. Watson, K.D.
 1688 Mansfield, Y. Pan, M.C. Simon, C.B. Thompson, and E. Gottlieb, Succinate
 1689 links TCA cycle dysfunction to oncogenesis by inhibiting HIF-alpha prolyl
 1690 hydroxylase. *Cancer Cell* 7 (2005) 77-85.

1691 M.I. Koukourakis, A. Giatromanolaki, E. Sivridis, K.C. Gatter, and A.L. Harris,
 1692 Pyruvate dehydrogenase and pyruvate dehydrogenase kinase expression in
 1693 non small cell lung cancer and tumor-associated stroma. *Neoplasia* 7 (2005)
 1694 1-6.

Legends to Figures

Figure 1. Viability and functionality of BCKDC-kinase knockdown *in vitro* system.

A, Western blot analysis showing sufficient knockdown of BDK corresponding to inhibition of phosphorylation and increased activity of BCKDC. C6 wild type (w/t), shK C6 and CVC6 cell lines are shown for the BCKDC-kinase, phospho-Ser292-E1 α BCKDC and total-E1 α BCKDC (n=6). Densitometric analyses of Western blots are depicted in densitometry units with standard errors normalized to wild type control. **B**, shows sufficient increased BCKDC activity resulting from decreased phosphorylation from BDK knockdown. Statistical differences were determined by Tukey's HSD *post hoc* analysis (*p<0.05, **p<0.001, and ***p<0.0001 with respect to C6 wild type; +p<0.05, ++p<0.001, and +++p<0.0001 with respect to CVC6; ND – non-detectable signal).

Figure 2. Analysis of BCKDC additional subunits

A, Representative Western blot analysis of additional BCKDC subunits, E1 β and E2, for C6 wild type (w/t), shK and CVC6 cells (n=6). **B**. Densitometric analyses of Western blots are depicted in densitometry units with standard errors normalized to wild type control. Statistical differences were determined by Tukey's HSD *post hoc* analysis (*p<0.05 with respect to C6 wild type).

Figure 3. Cell proliferation in serum and serum-free media show differences between shK cells and wild type and CVC6 control cells.

A, indicates cell proliferation is increased in shK cells with and without serum growth factors. Cell proliferation is shown in millions of cells for 24, 48, and 72 hr time points for viable cells only (n=6 for each time point). **B**. Cell proliferation in serum-free media is shown in millions of cells for 24, 48, and 72 hr time points for viable cells only (n=6 for each time point). Statistical differences were determined by Tukey's HSD *post hoc* analysis (*p<0.05 and ***p<0.0001 with respect to C6 wild type; +p<0.05, ++p<0.01, and +++p<0.0001 with respect to CVC6).

Figure 4. Metastatic phenotype analyses indicate differences between shK cells and wild type and CVC6 control cells.

A, shows increased migration for shK cells. Migrating C6 wild type(w/t), shK C6, and CVC6 cells were fixed and stained following 48 hour incubation (n=6). Representative images indicating qualitative differences are shown in **A** and quantitative cell counting is displayed in **D** with standard errors. **B**, shows increased invasion for shK cells. Invading C6 wild type(w/t), shK C6, and CVC6 cells were fixed and stained following 48 hour incubation (n=6). Representative images indicating qualitative differences are shown in **B** and quantitative cell counting is displayed in **E** with standard errors. **C**, shows increased colony diameter for shK cells indicating higher proliferation and invasion characteristics. Pictures of C6 wild type(w/t), shK C6, and CVC6 were taken following 3-week incubation in soft agar and analyzed for colony diameter. Representative images indicating qualitative differences are shown in **C** and quantitative colony diameter size is shown in **F** with standard errors. Statistical differences were determined Tukey's HSD *post hoc*

analyses (** $p < 0.0001$ with respect to C6 wild type; +++ $p < 0.0001$ with respect to CVC6).

Figure 5. Differential phosphorylation status of the mitochondrial metabolic complex PDC for shK and wild type and CVC6 controls


A, representative Western blot analysis shows phosphorylation status of Ser293-E1 α PDC for C6 wild type (w/t), shK C6 and CVC6 cells (n=6). **B**, indicates increased PDC phosphorylation ratios for shK cells. Densitometric analyses of Western blots are shown in densitometry units with standard errors normalized to wild type control. **C**, shows increased lactate levels for shK cells, which indicates an increased Warburg phenotype. Lactate production was measured from media by a CMA microdialysis analyzer at 6, 12, 24, and 48 hours (n=6 for each time point) normalized to cell number. Statistical differences were determined by Tukey's HSD *post hoc* analysis (* $p < 0.05$ with respect to C6 wild type; + $p < 0.05$ with respect to CVC6).

Table 1. Free ATP levels in C6 wild type, shK, and CVC6 cells

No significant difference between ATP levels was found. Data were gathered for shK, C6, and CVC6 cells by a luciferase bioluminescence assay. Data are displayed as nM ATP normalized by millions of viable cells (\pm SEM). Statistical analysis showed no significant difference between any of the cell lines (Tukey's HSD *post hoc* analysis, n=6).

1763 **Figures**


1764 **Figure 4. Viability and functionality of BCKDC-kinase knockdown in vitro**
1765 **system**



1766


1767

Figure 5. Analysis of BCKDC additional subunits



1768


1769 **Figure 6. Cell proliferation in serum and serum-free media show differences**
1770 **between shK cells and wild type and CVC6 control cells**



1771

1772

1773 **Figure 7. Metastatic phenotype analyses indicate differences between shK**
1774 **cells and wild type and CVC6 control cells**




1775

1776

1777

1778
1779

Figure 8. Differential phosphorylation status of the mitochondrial metabolic complex PDC for shK and wild type and CVC6 controls



1780
1781

1782
1783
1784
1785
1786

Table 1. Free ATP levels in C6 wild type, shK, and CVC6 cells	
Cell Type	nM ATP/million cells (means ± SEM)
C6 wild type	150 ± 23.4
shK C6	149 ± 2.5
CV C6	142 ± 7.3

General Discussion

The focus of this thesis is the role of BCAAs in cancer signaling and biochemistry. These data demonstrate a putative, novel signaling pathway involving HIF-1 α and that BCAA catabolism is not an independent regulator of *in vitro* cancer phenotypes. It has been known that the metabolism of cancer cells is aberrant in that they preferentially consume glucose anaerobically, even in the presence of oxygen levels sufficient for mitochondrial respiration. Recent findings from our lab have shown that manipulating PDC activity to be constitutively active (forcing aerobic respiration to occur) can reduce cancerous tumor formation and migration in a xenograph animal model. This work attempted to elaborate on the interest of another biochemical pathway, that of the BCAAs, in cancer biology. We set out to answer two research questions:

1. Does α -ketoisocaproic acid, a substrate of the BCKDC and deaminated product of leucine, act to decrease HIF degradation cycle activity thereby promoting neoplastic activity in an *in vitro* model of CNS cancer?
2. Do cells genetically modified by small hairpin RNA for the BCKDC kinase increase BCKDC activity and indicate a reciprocal relationship between BCKDC activity and cellular aggressiveness (eg, proliferation, migration, invasion, colony formation) in an *in vitro* model of CNS cancer, as similarly seen for PDC?

Chapter 1 focuses on the first question regarding BCKAs in HIF biology. It was first discovered in our lab that the BCKAs had a positive affect on HIF-1 α levels

in an *in vitro* system (Lu and others 2005). This work attempted to elaborate and further demonstrate that BCKAs can alter HIF regulation. Not only is HIF one of the most studied molecules in cancer biology, but, also, data have been shown that circulating BCAA levels are increased in patients with cancer. Increased circulating BCAA levels implies increased circulating BCKA levels, which, when taken together with our hypothesis for BCKA interference in HIF regulation, would have new implications in cancer biology.

Chapter 1, figure 1 shows dose-dependent curves for HIF-1 α protein and VEGF secretion in contrast to BCKDC phosphorylation status. It is clear from these data that BCKAs are able to affect HIF-1 α levels and VEGF secretion at relatively low levels, which fall in the range of BCAA levels found circulating in patients with cancer. Chapter 1, figure 2 shows BCKAs can inhibit the HPH cycle by sustaining HIF-1 α (endogenous HPH target) and ODD-GFP (exogenous reporter molecule of HPH cycle activity) levels. This figure also indicates that BCKA-induced HIF-1 α level stabilization is reversible. The mTOR molecule has been shown to be activated by leucine (Kimball and Jefferson 2004) and also to increase HIF-1 α levels (Hudson and others 2002). Chapter 1, figure 2 also shows increased HIF-1 α levels are not the result of mTOR activity.

Our research focused on the metabolism of BCAAs and BCKAs in two situations of potential clinical importance. The first involved the aberrant metabolism of BCKAs in maple syrup urine disease (MSUD), which leads us to propose ascorbate as a possible benign therapeutic intervention for treating MSUD-induced edema. Future work in this area could potentially extend these *in vitro* findings to a

physiological model of MSUD. The literature is still sparse with regards to mechanisms leading to the edematous phenotypes seen in MSUD patients in acute crisis. If the putative BCKA-HIF-VEGF pathway presented here is validated (either through *in vitro* models of MSUD or transgenic animals with inactive BCKDC), it may be beneficial to begin testing in MSUD patients. Ascorbate has long been known to either passively diffuse (Lam and Daniel 1986) or be transported via a specific Na-dependent transporter in the choroid plexus. Ascorbate levels have been shown to be 500 μ M in CSF (Stamford and others 1984), concentrated in the ventricular system (Spector and Lorenzo 1973). These levels are five-fold higher than necessary for activity *in vitro* (Knowles and others 2003).

The second project examined the effects of manipulation to BCAA metabolism on the resultant malignant phenotypes in cancer cell lines. While our initial aim was to identify BCKDC activation as a potential therapeutic mechanism, the research suggests that, at worst, the metabolic pathways of BCAAs are too intertwined with others to be clearly effective. In the best of circumstances, perhaps a line of research initiated by Doering and colleagues where BDK overexpression is used in an *in vitro* setting could indicate if artificially inactive BCKDC could lead to decreased malignant phenotypes, the inverse of what we have shown here.

The major finding shown in Chapter 2 was that a reduction of BDK activity, and thus an increase in BCKDC activity, increased *in vitro* aggressive and metastatic characteristics in C6 cells. This finding is surprising and the opposite of what was expected according to our rationale proposed where local depletion of

1855 essential amino acids would cause protein translation and *de novo* synthesis rates to
1856 decrease.

1857 Here we show short hairpin RNA interference successfully knocked down the
1858 translation of BDK. This kinase has been shown to phosphorylate Ser292 of the
1859 BCKDC E1 α subunit, causing deactivation (Harris and others 1997; Popov and
1860 others 1992; Shimomura and others 1990). This would allow the phosphatase to act
1861 unopposed, resulting in increased activation of BCKDC (Harris and others 2004).

1862 Through collaboration, the hypothesized increase in activation was verified through a
1863 BCKDC E1 α subunit activity assay. However, the increased activity was not to the
1864 proportions one would expect from the Western blots. Perhaps there is another
1865 kinase which phosphorylates a residue other than Ser292, leading to a decrease in
1866 BCKDC activity which would not be apparent in the Western blots. This seems
1867 unlikely due to the fact that various groups have consistently reported the isolation of
1868 one, and only one, kinase specific for BCKDC [reviewed in (Harris and others 2004;
1869 Harris and others 2005)]. There is another site phosphorylated by BDK (Ser³⁰²),
1870 although this has been shown to be a modification inconsequential to the activity of
1871 BCKDC (Cook and others 1984; Cook and others 1983; Li and others 2007). A more
1872 likely explanation is that, over time, the transfected cells have either rejected the
1873 plasmid, or the plasmid has mutated since the activity assays were the last
1874 chronological experiments performed. Both of these situations would result in the
1875 observed discrepancy, however, the latter is more likely due to the fact that the shK
1876 cells retained resistance to G418, the selection agent. A large body of evidence has
1877 shown tumor phenotypes such as proliferation, migration, invasion, and colony

formation are indicators of metastatic and aggressive cancers (Gao and others 2005; Ke and Costa 2006; Le Jeune and others 2006; Lin and others 2007; Martens and others 2006; Mohyeldin and others 2005; Mueller and others 1999; Zhang and others 2000). We performed experiments to analyze these characteristics to discover the effects of BDK knockdown on *in vitro* cancer phenotypes. Our hypothesis was that a reduction in BDK would increase BCKDC activity.

Consequently this would act

to deplete free BCAA from the pools necessary for other pathways, such as translational activation and anabolism for building new proteins *de novo*. The results of experiments testing these four metastatic phenotypes of our *in vitro*, transgenic GBM model

showed increased metastatic phenotypes, the opposite of what was expected. A summary of our

expected and actual results is shown in Fig 9. Interestingly, Nakai *et al.* showed a decrease in proliferation rates in C2C12 cells using transient transfection of small




Figure 9. Hypothesis and results summary for BDK knockdown experiments.

Diagram indicates our experimental objective to generate a BDK knockdown with our hypothesized results (left arm) and our experimental results (right arm). In the right arm, results obtained in accordance with our original hypothesis are encased in green lines; results counter to our original hypothesis are encased in red lines.

1901 inhibitory RNA to knockdown BDK. There are distinct differences in the Nakai *et al*,
1902 methods and those used in the present work. For instance, their use of transient
1903 transfection may have resulted in a smaller population of cells contain the
1904 knockdown plasmid. This would provide results that are representative of a mixed
1905 population of BDK expressing cells (Nakai and others 2006). The authors also
1906 indicated that cells were used for assays at 50% confluency. It is unclear from the
1907 methods whether they began the proliferation assays at such high confluency, but
1908 was probably necessary to achieve maximal transfection efficiency. Our proliferation
1909 assays began at closer to 25% confluency, allowing for more growth space and less
1910 inhibitory autoregulation of cellular growth. Also, differences in C2C12 and C6 cell
1911 phenotypes have not been defined, which may add to complications in the
1912 seemingly conflicting data. C2C12 are myogenic cells, whereas C6 are astrocyte-
1913 derived glioma cells. It is unclear what differences there are, if any, in the protein to
1914 lipid ratios in these cell types, which may yield variant results due to the unique
1915 aspects of BCAA metabolism discussed throughout this work. In addition, Nakai *et al*
1916 showed increased insulin secretion was sustained in the BDK siRNA-treated cells.
1917 Insulin has long been known to increase cellular viability and proliferation. Recent
1918 reports (Akhtar and others 2009; Kai and others 2009; Probst and others 2008) have
1919 shown that inhibiting the activity of insulin and insulin-like growth factors lead to
1920 decreased proliferation of cancerous cells. Therefore it seems plausible that there
1921 are other unidentified mechanisms at work in the differences in cellular proliferation
1922 showed by Nakai *et al* and the present work.

1923 Our second hypothesis was based on research performed in our laboratory
1924 where altering another mitochondrial dehydrogenase, PDC, reduced *in vitro*
1925 proliferation, invasion, migration, and colony formation (McFate and others 2008).
1926 Previous work had found that altering the phosphorylation status of Ser293-E1 α
1927 PDC (site 1) had effects analogous to what was originally hypothesized in this study,
1928 i.e. increasing PDC activation by interfering with the expression of pyruvate
1929 dehydrogenase complex kinase (PDK) decreased the metastatic phenotypes (Kim
1930 and others 2007; Koukourakis and others 2005; McFate and others 2008; Roche
1931 and Hiromasa 2007). The original research concerning PDC and cancer is related to
1932 the “Warburg Effect”, established in 1960 (Warburg 1930; Warburg 1956), and the
1933 “Pasteur Effect”, established in 1857 (Pasteur 1857; Pasteur 1859) [reviewed in
1934 English (Racker 1974)]. Warburg found that cancerous cells preferred glycolytic
1935 metabolism over cellular respiration, even in the presence of oxygen. His theory fell
1936 out of favor for 60 years until recently it has been revisited by researchers exploring
1937 the possibility that the metabolic changes actually cause cancer. Pasteur had
1938 focused more on the production of lactic acid as a byproduct of glycolytic
1939 metabolism in bacteria. These data combined formed the basis of PDC-linked
1940 cancer theories reported in the literature.

1941 We investigated whether there was a change in the phosphorylation state of
1942 E1 α PDC in shK cells, which could possibly explain the unforeseen increased
1943 metastatic phenotypes. We found a higher phosphorylation ratio of phosphorylated
1944 to total E1 α PDC for shK cells with respect to both wild type and CVC6 control cells
1945 typical of increased metastatic tendencies (Koukourakis and others 2005; McFate

and others 2008; Roche and Hiromasa 2007). When we analyzed “the Warburg Effect” by lactate analysis, we found that lactate production was increased for the shK cells compared to both wild type and CVC6 cells. These data compliment our previous data showing that PDC has more phosphorylation in shK cells. With PDC activity reduced, the cells accumulate pyruvate by conversion to lactate. These data showed that shK cells had decreased BDK levels, which may have caused a permanent shift in metabolism from PDC to the BCKDC in those cells.

New evidence has shown that “the Warburg Effect” may not be entirely about glucose metabolism. DeBerardinis *et al*, showed that SF188 glioma cells grown in culture consumed glutamine beyond levels required to accommodate nitrogen demand. In fact, the authors found glutamine metabolism primarily provided anaplerotic metabolites to the TCA cycle. The driving factors for this were the requirements of NADPH and TCA cycle intermediates used in fatty acid synthesis (DeBerardinis and others 2007). Since BCAA catabolism through BCKDC produces acetyl-CoA and other CoA derivatives, it is possible that fatty acid synthesis is also increased in the shK cells as a result of increased BCKDC activation through BDK knockdown. We focused, perhaps too narrowly, on the requirements of BCAAs for the *de novo* synthesis of proteins. However, dividing cells also require lipids to encase daughter cells after mitosis is complete. The relevant lipid/protein ratios are not clearly defined, however, it is known that mitochondrial inner membranes and pig epidermal cells have a protein to lipid ratio of 3:2, human red blood cell membranes and mouse liver ratios are 1:1, and myelin-producing cells have ratios closer to 1:5. These are just some examples of the variation of protein to lipid ratios found in

tissues. Joshi et al, report on BDK knockout mice that organ tissues have differential growth rates in these animals. Particularly, BDK knockout mice had decreased brain, muscle, and adipose tissue weights, similar heart weights, and increased liver and kidney weights when compared to wild type animals with respect to wild type. The authors posit the differential endogenous activity of BCKDC in these tissues for the inconsistent differences in tissue weight (Joshi and others 2007). For instance, the liver is known to have high endogenous BCKDC activity in wild-type mice. Therefore knocking out the BDK would not alter the metabolic pathways that may be more highly regulated in other tissues (eg, brain) so that the natural metabolic network is not shifted in the opposite direction, but rather in an accelerated manner.

To indulge in further speculation, a recent report by Singh et al, studying *Staphylococcus aureus* where the investigators disrupted the transcription of BCKDC-akin genes in the bacterial genome led to decreased branched-chain fatty acid production. This alteration led to increased susceptibility to stresses and, therefore, decreased the viability of the cells (Singh and others 2008). This research agrees with another recent report showing inhibition of LAT1 transporters in eukaryotic cancer cells decreases cell proliferation and increases apoptosis through caspase 3 and 7 activities. These data indicate that shutting down the availability of BCAAs to the metabolic milieu of the cell decreases the viability of the cell. We elected to investigate if the role of the BCKDC could affect proliferation rates in a similar vane. Our results are contrary to our hypothesis about decreasing the availability of BCAAs for anabolic pathways, but it does seem that limiting BCAA entry into the metabolic milieu upstream of the BCKDC-involved processes does

result in decreased proliferation. Perhaps further analyses on branched-chain fatty acid production, carbon skeleton tracking, and anabolic protein turnover would provide evidence to explain the unforeseen phenotypes seen in the shK cells and how alternate compensatory biochemical changes seem to have occurred.

GBMs are difficult to treat, partially due to their location in the highly organized and compacted brain tissue. However, since these cells are hypermetabolic, it is possible that they may be preferentially susceptible to selective metabolic attack. The goal of this research was to investigate the potential of the BCKDC to be such a therapeutic target. Advances in targeted strategies for brain cancers have shown some interesting and promising results [reviewed in (Fine 2007; Lukas and others 2007)]. However, a major drawback of targeted therapies is their specificity for cancers with narrowly defined etiologies. Our strategy was an attempt to identify a more global target for cancer therapeutic agents. We hoped to find that BCKDC would be a potential target for further research and perhaps rational drug design for new, global cancer treatments. In light of the research supporting that cancer cachexia is the result of increased BCAA uptake in cancerous tissues, causing catabolic breakdown of skeletal muscle, we hypothesized that BCKDC would indeed lead to a novel therapeutic target. However, the data presented here indicate there is more to the story than BCAA shunting, but rather that this biochemical pathway may be regulated by other pathways that effect cancer cell phenotypes.

It became apparent in the early stages of this research that the area of BCAA involvement in metabolic and signaling pathways pertinent to cancer biology lacked

2015 a seminal work to bring a comprehensive direction and strategy to research in this
2016 field. The rationale for the hypotheses which lay the foundation for this thesis was
2017 composed of research investigations in exercise physiology, bacterial and
2018 mammalian biochemistry, cancer cell metabolism, cancer's impact on muscular
2019 physiology and biochemistry, and genetic disorders of BCAA metabolism, to name a
2020 few. Recent reports discussed in this section do, however, begin examining the
2021 impact of BCAA metabolic changes in cancer cells. It is my opinion that this field of
2022 research has been building momentum and may be an important topic in cancer
2023 research in the near future.

2024 **References**

- 2025 Akhtar S, Meeran SM, Katiyar N, Katiyar SK. 2009. Grape seed proanthocyanidins
2026 inhibit the growth of human non-small cell lung cancer xenografts by targeting
2027 insulin-like growth factor binding protein-3, tumor cell proliferation, and
2028 angiogenic factors. *Clin Cancer Res* 15(3):821-31.
- 2029 Aldape KD, Okcu MF, Bondy ML, Wrensch M. 2003. Molecular epidemiology of
2030 glioblastoma. *Cancer J* 9(2):99-106.
- 2031 Anthony JC, Anthony TG, Kimball SR, Vary TC, Jefferson LS. 2000a. Orally
2032 administered leucine stimulates protein synthesis in skeletal muscle of
2033 postabsorptive rats in association with increased eIF4F formation. *J Nutr*
2034 130(2):139-45.
- 2035 Anthony JC, Yoshizawa F, Anthony TG, Vary TC, Jefferson LS, Kimball SR. 2000b.
2036 Leucine stimulates translation initiation in skeletal muscle of postabsorptive
2037 rats via a rapamycin-sensitive pathway. *J Nutr* 130(10):2413-9.
- 2038 Arany Z, Huang LE, Eckner R, Bhattacharya S, Jiang C, Goldberg MA, Bunn HF,
2039 Livingston DM. 1996. An essential role for p300/CBP in the cellular response
2040 to hypoxia. *Proc Natl Acad Sci U S A* 93(23):12969-73.
- 2041 Armstrong RC. 1998. Isolation and characterization of immature oligodendrocyte
2042 lineage cells. *Methods* 16(3):282-92.
- 2043 Baracos VE. 2000. Regulation of skeletal-muscle-protein turnover in cancer-
2044 associated cachexia. *Nutrition* 16(10):1015-8.
- 2045 Baracos VE, Mackenzie ML. 2006. Investigations of branched-chain amino acids
2046 and their metabolites in animal models of cancer. *J Nutr* 136(1 Suppl):237S-
2047 42S.
- 2048 Benda P, Lightbody J, Sato G, Levine L, Sweet W. 1968. Differentiated rat glial cell
2049 strain in tissue culture. *Science* 161(839):370-1.
- 2050 Bixel MG, Hutson SM, Hamprecht B. 1997. Cellular distribution of branched-chain
2051 amino acid aminotransferase isoenzymes among rat brain glial cells in
2052 culture. *J Histochem Cytochem* 45(5):685-94.
- 2053 Brismar J, Aqeel A, Brismar G, Coates R, Gascon G, Ozand P. 1990. Maple syrup
2054 urine disease: findings on CT and MR scans of the brain in 10 infants. *AJNR*
2055 *Am J Neuroradiol* 11(6):1219-28.
- 2056 Brown EJ, Albers MW, Shin TB, Ichikawa K, Keith CT, Lane WS, Schreiber SL.
2057 1994. A mammalian protein targeted by G1-arresting rapamycin-receptor
2058 complex. *Nature* 369(6483):756-8.

- 2059 Bruick RK, McKnight SL. 2001. A conserved family of prolyl-4-hydroxylases that
2060 modify HIF. *Science* 294(5545):1337-40.
- 2061 Brunn GJ, Hudson CC, Sekulic A, Williams JM, Hosoi H, Houghton PJ, Lawrence
2062 JC, Jr., Abraham RT. 1997. Phosphorylation of the translational repressor
2063 PHAS-I by the mammalian target of rapamycin. *Science* 277(5322):99-101.
- 2064 Burton EC, Prados MD. 2000. Malignant gliomas. *Curr Treat Options Oncol*
2065 1(5):459-68.
- 2066 Chen C, Pore N, Behrooz A, Ismail-Beigi F, Maity A. 2001. Regulation of glut1
2067 mRNA by hypoxia-inducible factor-1. Interaction between H-ras and hypoxia.
2068 *J Biol Chem* 276(12):9519-25.
- 2069 Chuang DT, Chuang JL, Wynn RM. 2006. Lessons from genetic disorders of
2070 branched-chain amino acid metabolism. *J Nutr* 136(1 Suppl):243S-9S.
- 2071 Cook KG, Bradford AP, Yeaman SJ, Aitken A, Fearnley IM, Walker JE. 1984.
2072 Regulation of bovine kidney branched-chain 2-oxoacid dehydrogenase
2073 complex by reversible phosphorylation. *Eur J Biochem* 145(3):587-91.
- 2074 Cook KG, Lawson R, Yeaman SJ. 1983. Multi-site phosphorylation of bovine kidney
2075 branched-chain 2-oxoacid dehydrogenase complex. *FEBS Lett* 157(1):59-62.
- 2076 Costelli P, Llovera M, Garcia-Martinez C, Carbo N, Lopez-Soriano FJ, Argiles JM.
2077 1995. Enhanced leucine oxidation in rats bearing an ascites hepatoma
2078 (Yoshida AH-130) and its reversal by clenbuterol. *Cancer Lett* 91(1):73-8.
- 2079 Cota D, Proulx K, Smith KA, Kozma SC, Thomas G, Woods SC, Seeley RJ. 2006.
2080 Hypothalamic mTOR signaling regulates food intake. *Science* 312(5775):927-
2081 30.
- 2082 D'Angelo G, Duplan E, Vigne P, Frelin C. 2003. Cyclosporin A prevents the hypoxic
2083 adaptation by activating hypoxia-inducible factor-1alpha Pro-564
2084 hydroxylation. *J Biol Chem* 278(17):15406-11.
- 2085 Dalgard CL, Lu H, Mohyeldin A, Verma A. 2004. Endogenous 2-oxoacids
2086 differentially regulate expression of oxygen sensors. *Biochem J* 380(Pt
2087 2):419-24.
- 2088 Damuni Z, Merryfield ML, Humphreys JS, Reed LJ. 1984. Purification and properties
2089 of branched-chain alpha-keto acid dehydrogenase phosphatase from bovine
2090 kidney. *Proc Natl Acad Sci U S A* 81(14):4335-8.
- 2091 Damuni Z, Reed LJ. 1987. Purification and properties of the catalytic subunit of the
2092 branched-chain alpha-keto acid dehydrogenase phosphatase from bovine
2093 kidney mitochondria. *J Biol Chem* 262(11):5129-32.

- 2094 Dancis J, Levitz M, Westall RG. 1960. Maple syrup urine disease: branched-chain
2095 keto-aciduria. *Pediatrics* 25:72-9.
- 2096 Danner DJ, Lemmon SK, Elsas LJ, 2nd. 1978. Substrate specificity and stabilization
2097 by thiamine pyrophosphate of rat liver branched chain alpha-ketoacid
2098 dehydrogenase. *Biochem Med* 19(1):27-38.
- 2099 DeBerardinis RJ, Mancuso A, Daikhin E, Nissim I, Yudkoff M, Wehrli S, Thompson
2100 CB. 2007. Beyond aerobic glycolysis: transformed cells can engage in
2101 glutamine metabolism that exceeds the requirement for protein and
2102 nucleotide synthesis. *Proc Natl Acad Sci U S A* 104(49):19345-50.
- 2103 Denko NC, Fontana LA, Hudson KM, Sutphin PD, Raychaudhuri S, Altman R,
2104 Giaccia AJ. 2003. Investigating hypoxic tumor physiology through gene
2105 expression patterns. *Oncogene* 22(37):5907-14.
- 2106 Erecinska M, Nelson D, Wilson DF, Silver IA. 1984. Neurotransmitter amino acids in
2107 the CNS. I. Regional changes in amino acid levels in rat brain during ischemia
2108 and reperfusion. *Brain Res* 304(1):9-22.
- 2109 Fine HA. 2007. Promising new therapies for malignant gliomas. *Cancer J* 13(6):349-
2110 54.
- 2111 Gao CF, Xie Q, Su YL, Koeman J, Khoo SK, Gustafson M, Knudsen BS, Hay R,
2112 Shinomiya N, Vande Woude GF. 2005. Proliferation and invasion: plasticity in
2113 tumor cells. *Proc Natl Acad Sci U S A* 102(30):10528-33.
- 2114 Goldberg MA, Dunning SP, Bunn HF. 1988. Regulation of the erythropoietin gene:
2115 evidence that the oxygen sensor is a heme protein. *Science* 242(4884):1412-
2116 5.
- 2117 Gradin K, McGuire J, Wenger RH, Kvietikova I, fhitelaw ML, Toftgard R, Tora L,
2118 Gassmann M, Poellinger L. 1996. Functional interference between hypoxia
2119 and dioxin signal transduction pathways: competition for recruitment of the
2120 Arnt transcription factor. *Mol Cell Biol* 16(10):5221-31.
- 2121 Graven KK, Yu Q, Pan D, Roncarati JS, Farber HW. 1999. Identification of an
2122 oxygen responsive enhancer element in the glyceraldehyde-3-phosphate
2123 dehydrogenase gene. *Biochim Biophys Acta* 1447(2-3):208-18.
- 2124 Greenberg R, Reaven G. 1966. The effect of L-leucine on hepatic glucose formation.
2125 *Pediatrics* 37(6):934-41.
- 2126 Grobбен B, De Deyn PP, Slegers H. 2002. Rat C6 glioma as experimental model
2127 system for the study of glioblastoma growth and invasion. *Cell Tissue Res*
2128 310(3):257-70.

- 2129 Hall TR, Wallin R, Reinhart GD, Hutson SM. 1993. Branched chain
2130 aminotransferase isoenzymes. Purification and characterization of the rat
2131 brain isoenzyme. *J Biol Chem* 268(5):3092-8.
- 2132 Hanahan D, Weinberg RA. 2000. The hallmarks of cancer. *Cell* 100(1):57-70.
- 2133 Harper AE. 1989. Thoughts on the role of branched-chain alpha-keto acid
2134 dehydrogenase complex in nitrogen metabolism. *Ann N Y Acad Sci* 573:267-
2135 73.
- 2136 Harper AE, Miller RH, Block KP. 1984. Branched-chain amino acid metabolism.
2137 *Annu Rev Nutr* 4:409-54.
- 2138 Harris RA, Hawes JW, Popov KM, Zhao Y, Shimomura Y, Sato J, Jaskiewicz J,
2139 Hurley TD. 1997. Studies on the regulation of the mitochondrial alpha-
2140 ketoacid dehydrogenase complexes and their kinases. *Adv Enzyme Regul*
2141 37:271-93.
- 2142 Harris RA, Joshi M, Jeoung NH. 2004. Mechanisms responsible for regulation of
2143 branched-chain amino acid catabolism. *Biochem Biophys Res Commun*
2144 313(2):391-6.
- 2145 Harris RA, Joshi M, Jeoung NH, Obayashi M. 2005. Overview of the molecular and
2146 biochemical basis of branched-chain amino acid catabolism. *J Nutr* 135(6
2147 Suppl):1527S-30S.
- 2148 Harris RA, Popov KM, Zhao Y, Kedishvili NY, Shimomura Y, Crabb DW. 1995. A
2149 new family of protein kinases--the mitochondrial protein kinases. *Adv Enzyme*
2150 *Regul* 35:147-62.
- 2151 Hewitson KS, McNeill LA, Riordan MV, Tian YM, Bullock AN, Welford RW, Elkins
2152 JM, Oldham NJ, Bhattacharya S, Gleadle JM and others. 2002. Hypoxia-
2153 inducible factor (HIF) asparagine hydroxylase is identical to factor inhibiting
2154 HIF (FIH) and is related to the cupin structural family. *J Biol Chem*
2155 277(29):26351-5.
- 2156 Hon WC, Wilson MI, Harlos K, Claridge TD, Schofield CJ, Pugh CW, Maxwell PH,
2157 Ratcliffe PJ, Stuart DI, Jones EY. 2002. Structural basis for the recognition of
2158 hydroxyproline in HIF-1 alpha by pVHL. *Nature* 417(6892):975-8.
- 2159 Hudson CC, Liu M, Chiang GG, Otterness DM, Loomis DC, Kaper F, Giaccia AJ,
2160 Abraham RT. 2002. Regulation of hypoxia-inducible factor 1alpha expression
2161 and function by the mammalian target of rapamycin. *Mol Cell Biol*
2162 22(20):7004-14.
- 2163 Humar R, Kiefer FN, Berns H, Resink TJ, Battegay EJ. 2002. Hypoxia enhances
2164 vascular cell proliferation and angiogenesis in vitro via rapamycin (mTOR)-
2165 dependent signaling. *FASEB J* 16(8):771-80.

- 2166 Hunter DC, Weintraub M, Blackburn GL, Bistrian BR. 1989. Branched chain amino
2167 acids as the protein component of parenteral nutrition in cancer cachexia. *Br*
2168 *J Surg* 76(2):149-53.
- 2169 Hutson SM, Fenstermacher D, Mahar C. 1988. Role of mitochondrial transamination
2170 in branched chain amino acid metabolism. *J Biol Chem* 263(8):3618-25.
- 2171 Hutson SM, Hall TR. 1993. Identification of the mitochondrial branched chain
2172 aminotransferase as a branched chain alpha-keto acid transport protein. *J*
2173 *Biol Chem* 268(5):3084-91.
- 2174 Hutson SM, Wallin R, Hall TR. 1992. Identification of mitochondrial branched chain
2175 aminotransferase and its isoforms in rat tissues. *J Biol Chem* 267(22):15681-
2176 6.
- 2177 Inculet RI, Stein TP, Peacock JL, Leskiw M, Maher M, Gorschboth CM, Norton JA.
2178 1987. Altered leucine metabolism in noncachectic sarcoma patients. *Cancer*
2179 *Res* 47(17):4746-9.
- 2180 Inui A. 2002. Cancer anorexia-cachexia syndrome: current issues in research and
2181 management. *CA Cancer J Clin* 52(2):72-91.
- 2182 Isaacs JS, Jung YJ, Mole DR, Lee S, Torres-Cabala C, Chung YL, Merino M, Trepel
2183 J, Zbar B, Toro J and others. 2005. HIF overexpression correlates with
2184 biallelic loss of fumarate hydratase in renal cancer: novel role of fumarate in
2185 regulation of HIF stability. *Cancer Cell* 8(2):143-53.
- 2186 Ivan M, Kondo K, Yang H, Kim W, Valiando J, Ohh M, Salic A, Asara JM, Lane WS,
2187 Kaelin WG, Jr. 2001. HIFalpha targeted for VHL-mediated destruction by
2188 proline hydroxylation: implications for O2 sensing. *Science* 292(5516):464-8.
- 2189 Jaakkola P, Mole DR, Tian YM, Wilson MI, Gielbert J, Gaskell SJ, Kriegsheim A,
2190 Hebestreit HF, Mukherji M, Schofield CJ and others. 2001. Targeting of HIF-
2191 alpha to the von Hippel-Lindau ubiquitylation complex by O2-regulated prolyl
2192 hydroxylation. *Science* 292(5516):468-72.
- 2193 Jeoung NH, Wu P, Joshi MA, Jaskiewicz J, Bock CB, Depaoli-Roach AA, Harris RA.
2194 2006. Role of pyruvate dehydrogenase kinase isoenzyme 4 (PDHK4) in
2195 glucose homeostasis during starvation. *Biochem J* 397(3):417-25.
- 2196 Joshi M, Jeoung NH, Popov KM, Harris RA. 2007. Identification of a novel PP2C-
2197 type mitochondrial phosphatase. *Biochem Biophys Res Commun* 356(1):38-
2198 44.
- 2199 Josko J, Gwozdz B, Jedrzejowska-Szypulka H, Hendryk S. 2000. Vascular
2200 endothelial growth factor (VEGF) and its effect on angiogenesis. *Med Sci*
2201 *Monit* 6(5):1047-52.

- 2202 Kai K, D'Costa S, Sills RC, Kim Y. 2009. Inhibition of the insulin-like growth factor 1
2203 receptor pathway enhances the antitumor effect of cisplatin in human
2204 malignant mesothelioma cell lines. *Cancer Lett*.
- 2205 Kallio PJ, Pongratz I, Gradin K, McGuire J, Poellinger L. 1997. Activation of hypoxia-
2206 inducible factor 1alpha: posttranscriptional regulation and conformational
2207 change by recruitment of the Arnt transcription factor. *Proc Natl Acad Sci U S*
2208 *A* 94(11):5667-72.
- 2209 Kanamori K, Ross BD, Kondrat RW. 1998. Rate of glutamate synthesis from leucine
2210 in rat brain measured in vivo by ¹⁵N NMR. *J Neurochem* 70(3):1304-15.
- 2211 Ke Q, Costa M. 2006. Hypoxia-inducible factor-1 (HIF-1). *Mol Pharmacol*
2212 70(5):1469-80.
- 2213 Kim JW, Gao P, Liu YC, Semenza GL, Dang CV. 2007. Hypoxia-Inducible Factor 1
2214 and Dysregulated c-Myc Cooperatively Induce Vascular Endothelial Growth
2215 Factor and Metabolic Switches Hexokinase 2 and Pyruvate Dehydrogenase
2216 Kinase 1. *Mol Cell Biol* 27(21):7381-93.
- 2217 Kim JW, Tchernyshyov I, Semenza GL, Dang CV. 2006. HIF-1-mediated expression
2218 of pyruvate dehydrogenase kinase: a metabolic switch required for cellular
2219 adaptation to hypoxia. *Cell Metab* 3(3):177-85.
- 2220 Kimball SR, Jefferson LS. 2004. Regulation of global and specific mRNA translation
2221 by oral administration of branched-chain amino acids. *Biochem Biophys Res*
2222 *Commun* 313(2):423-7.
- 2223 Kimball SR, Jefferson LS. 2006. New functions for amino acids: effects on gene
2224 transcription and translation. *Am J Clin Nutr* 83(2):500S-507S.
- 2225 Kimball SR, Jefferson LS, Fadden P, Haystead TA, Lawrence JC, Jr. 1996. Insulin
2226 and diabetes cause reciprocal changes in the association of eIF-4E and
2227 PHAS-I in rat skeletal muscle. *Am J Physiol* 270(2 Pt 1):C705-9.
- 2228 Kimball SR, Shantz LM, Horetsky RL, Jefferson LS. 1999. Leucine regulates
2229 translation of specific mRNAs in L6 myoblasts through mTOR-mediated
2230 changes in availability of eIF4E and phosphorylation of ribosomal protein S6.
2231 *J Biol Chem* 274(17):11647-52.
- 2232 Knowles HJ, Raval RR, Harris AL, Ratcliffe PJ. 2003. Effect of ascorbate on the
2233 activity of hypoxia-inducible factor in cancer cells. *Cancer Res* 63(8):1764-8.
- 2234 Koukourakis MI, Giatromanolaki A, Sivridis E, Gatter KC, Harris AL. 2005. Pyruvate
2235 dehydrogenase and pyruvate dehydrogenase kinase expression in non small
2236 cell lung cancer and tumor-associated stroma. *Neoplasia* 7(1):1-6.

- 2237 Lam DK, Daniel PM. 1986. The influx of ascorbic acid into the rat's brain. *Q J Exp*
2238 *Physiol* 71(3):483-9.
- 2239 Lam TK, Gutierrez-Juarez R, Pocai A, Rossetti L. 2005. Regulation of blood glucose
2240 by hypothalamic pyruvate metabolism. *Science* 309(5736):943-7.
- 2241 Lambert DG, Hughes K, Atkins TW. 1986. Insulin release from a cloned hamster B-
2242 cell line (HIT-T15). The effects of glucose, amino acids, sulphonylureas and
2243 colchicine. *Biochem Biophys Res Commun* 140(2):616-25.
- 2244 Lando D, Peet DJ, Gorman JJ, Whelan DA, Whitelaw ML, Bruick RK. 2002a. FIH-1
2245 is an asparaginyl hydroxylase enzyme that regulates the transcriptional
2246 activity of hypoxia-inducible factor. *Genes Dev* 16(12):1466-71.
- 2247 Lando D, Peet DJ, Whelan DA, Gorman JJ, Whitelaw ML. 2002b. Asparagine
2248 hydroxylation of the HIF transactivation domain a hypoxic switch. *Science*
2249 295(5556):858-61.
- 2250 Le Jeune N, Perek N, Dubois F. 2006. Influence of Pi3-K and PKC activity on
2251 99mTc-(V)-DMSA uptake: correlation with tumour aggressiveness in an in
2252 vitro malignant glioblastoma cell line model. *Eur J Nucl Med Mol Imaging*
2253 33(10):1206-13.
- 2254 Levy AP, Levy NS, Wegner S, Goldberg MA. 1995. Transcriptional regulation of the
2255 rat vascular endothelial growth factor gene by hypoxia. *J Biol Chem*
2256 270(22):13333-40.
- 2257 Li J, Machius M, Chuang JL, Wynn RM, Chuang DT. 2007. The two active sites in
2258 human branched-chain alpha-keto acid dehydrogenase operate
2259 independently without an obligatory alternating-site mechanism. *J Biol Chem*
2260 282(16):11904-13.
- 2261 Lin J, Chen L, Lin Z, Zhao M. 2007. Inhibitory effect of triptolide on glioblastoma
2262 multiforme in vitro. *J Int Med Res* 35(4):490-6.
- 2263 Long W, Saffer L, Wei L, Barrett EJ. 2000. Amino acids regulate skeletal muscle
2264 PHAS-I and p70 S6-kinase phosphorylation independently of insulin. *Am J*
2265 *Physiol Endocrinol Metab* 279(2):E301-6.
- 2266 Lorite MJ, Cariuk P, Tisdale MJ. 1997. Induction of muscle protein degradation by a
2267 tumour factor. *Br J Cancer* 76(8):1035-40.
- 2268 Lu H, Dalgard CL, Mohyeldin A, McFate T, Tait AS, Verma A. 2005. Reversible
2269 inactivation of HIF-1 prolyl hydroxylases allows cell metabolism to control
2270 basal HIF-1. *J Biol Chem* 280(51):41928-39.

- 2271 Lu H, Forbes RA, Verma A. 2002. Hypoxia-inducible factor 1 activation by aerobic
2272 glycolysis implicates the Warburg effect in carcinogenesis. *J Biol Chem*
2273 277(26):23111-5.
- 2274 Lukas RV, Boire A, Nicholas MK. 2007. Emerging therapies for malignant glioma.
2275 *Expert Rev Anticancer Ther* 7(12 Suppl):S29-36.
- 2276 Lundholm K, Bylund AC, Holm J, Schersten T. 1976. Skeletal muscle metabolism in
2277 patients with malignant tumor. *Eur J Cancer* 12(6):465-73.
- 2278 Lynch CJ, Halle B, Fujii H, Vary TC, Wallin R, Damuni Z, Hutson SM. 2003. Potential
2279 role of leucine metabolism in the leucine-signaling pathway involving mTOR.
2280 *Am J Physiol Endocrinol Metab* 285(4):E854-63.
- 2281 MacDonald MJ, Fahien LA, Mertz RJ, Rana RS. 1989. Effect of esters of succinic
2282 acid and other citric acid cycle intermediates on insulin release and inositol
2283 phosphate formation by pancreatic islets. *Arch Biochem Biophys* 269(2):400-
2284 6.
- 2285 Manalo DJ, Rowan A, Lavoie T, Natarajan L, Kelly BD, Ye SQ, Garcia JG, Semenza
2286 GL. 2005. Transcriptional regulation of vascular endothelial cell responses to
2287 hypoxia by HIF-1. *Blood* 105(2):659-69.
- 2288 Martens T, Schmidt NO, Eckerich C, Fillbrandt R, Merchant M, Schwall R, Westphal
2289 M, Lamszus K. 2006. A novel one-armed anti-c-Met antibody inhibits
2290 glioblastoma growth in vivo. *Clin Cancer Res* 12(20 Pt 1):6144-52.
- 2291 Masson N, Willam C, Maxwell PH, Pugh CW, Ratcliffe PJ. 2001. Independent
2292 function of two destruction domains in hypoxia-inducible factor-alpha chains
2293 activated by prolyl hydroxylation. *EMBO J* 20(18):5197-206.
- 2294 Mathupala SP, Rempel A, Pedersen PL. 2001. Glucose catabolism in cancer cells:
2295 identification and characterization of a marked activation response of the type
2296 II hexokinase gene to hypoxic conditions. *J Biol Chem* 276(46):43407-12.
- 2297 McFate T, Mohyeldin A, Lu H, Thakar J, Henriques J, Halim ND, Wu H, Schell MJ,
2298 Tsang TM, Teahan O and others. 2008. Pyruvate dehydrogenase complex
2299 activity controls metabolic and malignant phenotype in cancer cells. *J Biol*
2300 *Chem* 283(33):22700-8.
- 2301 Medhkour A, Chan M. 2005. Extremely rare glioblastoma multiforme of the conus
2302 medullaris with holocord and brain stem metastases, leading to cranial nerve
2303 deficit and respiratory failure: a case report and review of the literature. *Surg*
2304 *Neurol* 63(6):576-82; discussion 582-3.
- 2305 Miller CR, Perry A. 2007. Glioblastoma. *Arch Pathol Lab Med* 131(3):397-406.

- 2306 Min JH, Yang H, Ivan M, Gertler F, Kaelin WG, Jr., Pavletich NP. 2002. Structure of
2307 an HIF-1alpha -pVHL complex: hydroxyproline recognition in signaling.
2308 Science 296(5574):1886-9.
- 2309 Mohyeldin A, Lu H, Dalgard C, Lai SY, Cohen N, Acs G, Verma A. 2005.
2310 Erythropoietin signaling promotes invasiveness of human head and neck
2311 squamous cell carcinoma. Neoplasia 7(5):537-43.
- 2312 Mordier S, Deval C, Bechet D, Tassa A, Ferrara M. 2000. Leucine limitation induces
2313 autophagy and activation of lysosome-dependent proteolysis in C2C12
2314 myotubes through a mammalian target of rapamycin-independent signaling
2315 pathway. J Biol Chem 275(38):29900-6.
- 2316 Mueller MM, Herold-Mende CC, Riede D, Lange M, Steiner HH, Fusenig NE. 1999.
2317 Autocrine growth regulation by granulocyte colony-stimulating factor and
2318 granulocyte macrophage colony-stimulating factor in human gliomas with
2319 tumor progression. Am J Pathol 155(5):1557-67.
- 2320 Mujic A, Hunn A, Taylor AB, Lowenthal RM. 2006. Extracranial metastases of a
2321 glioblastoma multiforme to the pleura, small bowel and pancreas. J Clin
2322 Neurosci 13(6):677-81.
- 2323 Murray B, Wilson DJ. 2001. A study of metabolites as intermediate effectors in
2324 angiogenesis. Angiogenesis 4(1):71-7.
- 2325 Nakai N, Shimomura Y, Tamura T, Tamura N, Hamada K, Kawano F, Ohira Y. 2006.
2326 Leucine-induced activation of translational initiation is partly regulated by the
2327 branched-chain alpha-keto acid dehydrogenase complex in C2C12 cells.
2328 Biochem Biophys Res Commun 343(4):1244-50.
- 2329 Newton HB, Rosenblum MK, Walker RW. 1992. Extraneural metastases of
2330 infratentorial glioblastoma multiforme to the peritoneal cavity. Cancer
2331 69(8):2149-53.
- 2332 Nieder C, Grosu AL, Astner S, Molls M. 2005. Treatment of unresectable
2333 glioblastoma multiforme. Anticancer Res 25(6C):4605-10.
- 2334 Noda C, Ichihara A. 1974. Control of ketogenesis from amino acids. II. Ketone
2335 bodies formation from alpha-ketoisocaproate, the keto-analogue of leucine,
2336 by rat liver mitochondria. J Biochem 76(5):1123-30.
- 2337 Noda C, Ichihara A. 1976. Control of ketogenesis from amino acids. IV. Tissue
2338 specificity in oxidation of leucine, tyrosine, and lysine. J Biochem 80(5):1159-
2339 64.
- 2340 O'Keefe SJ, Ogden J, Ramjee G, Rund J. 1990. Contribution of elevated protein
2341 turnover and anorexia to cachexia in patients with hepatocellular carcinoma.
2342 Cancer Res 50(4):1226-30.

- 2343 Ogawa K, Yokojima A, Ichihara A. 1970. Transaminase of branched chain amino
2344 acids. VII. Comparative studies on isozymes of ascites hepatoma and various
2345 normal tissues of rat. J Biochem 68(6):901-11.
- 2346 Papandreou I, Cairns RA, Fontana L, Lim AL, Denko NC. 2006. HIF-1 mediates
2347 adaptation to hypoxia by actively downregulating mitochondrial oxygen
2348 consumption. Cell Metab 3(3):187-97.
- 2349 Parker PJ, Randle PJ. 1978. Partial purification and properties of branched-chain 2-
2350 oxo acid dehydrogenase of ox liver. Biochem J 171(3):751-7.
- 2351 Pasteur L. 1857. Memoire sur la fermentation appelee lactique. Comptes rendus de
2352 l'Academie des sciences 45:3.
- 2353 Pasteur L. 1859. Nouveaux faits pour servir a l'histoire de la levure lactique.
2354 Comptes rendus de l'Academie des sciences 48:2.
- 2355 Pettit FH, Yeaman SJ, Reed LJ. 1978. Purification and characterization of branched
2356 chain alpha-keto acid dehydrogenase complex of bovine kidney. Proc Natl
2357 Acad Sci U S A 75(10):4881-5.
- 2358 Pollard PJ, Briere JJ, Alam NA, Barwell J, Barclay E, Wortham NC, Hunt T, Mitchell
2359 M, Olpin S, Moat SJ and others. 2005. Accumulation of Krebs cycle
2360 intermediates and over-expression of HIF1alpha in tumours which result from
2361 germline FH and SDH mutations. Hum Mol Genet 14(15):2231-9.
- 2362 Popov KM, Zhao Y, Shimomura Y, Kuntz MJ, Harris RA. 1992. Branched-chain
2363 alpha-ketoacid dehydrogenase kinase. Molecular cloning, expression, and
2364 sequence similarity with histidine protein kinases. J Biol Chem
2365 267(19):13127-30.
- 2366 Probst OC, Puxbaum V, Svoboda B, Leksa V, Stockinger H, Mikula M, Mikulits W,
2367 Mach L. 2008. The mannose 6-phosphate/insulin-like growth factor II receptor
2368 restricts the tumourigenicity and invasiveness of squamous cell carcinoma
2369 cells. Int J Cancer.
- 2370 Racker E. 1974. History of the Pasteur effect and its pathobiology. Mol Cell Biochem
2371 5(1-2):17-23.
- 2372 Riviello JJ, Jr., Rezvani I, DiGeorge AM, Foley CM. 1991. Cerebral edema causing
2373 death in children with maple syrup urine disease. J Pediatr 119(1 (Pt 1)):42-
2374 5.
- 2375 Roberts CM, Sokatch JR. 1978. Branched chain amino acids as activators of
2376 branched chain ketoacid dehydrogenase. Biochem Biophys Res Commun
2377 82(3):828-33.

- 2378 Robey IF, Lien AD, Welsh SJ, Baggett BK, Gillies RJ. 2005. Hypoxia-inducible
2379 factor-1alpha and the glycolytic phenotype in tumors. *Neoplasia* 7(4):324-30.
- 2380 Roche TE, Hiromasa Y. 2007. Pyruvate dehydrogenase kinase regulatory
2381 mechanisms and inhibition in treating diabetes, heart ischemia, and cancer.
2382 *Cell Mol Life Sci* 64(7-8):830-49.
- 2383 Saad AG, Sachs J, Turner CD, Proctor M, Marcus KJ, Wang L, Lidov H, Ullrich NJ.
2384 2007. Extracranial metastases of glioblastoma in a child: case report and
2385 review of the literature. *J Pediatr Hematol Oncol* 29(3):190-4.
- 2386 Sabatini DM, Erdjument-Bromage H, Lui M, Tempst P, Snyder SH. 1994. RAFT1: a
2387 mammalian protein that binds to FKBP12 in a rapamycin-dependent fashion
2388 and is homologous to yeast TORs. *Cell* 78(1):35-43.
- 2389 Sabers CJ, Martin MM, Brunn GJ, Williams JM, Dumont FJ, Wiederrecht G,
2390 Abraham RT. 1995. Isolation of a protein target of the FKBP12-rapamycin
2391 complex in mammalian cells. *J Biol Chem* 270(2):815-22.
- 2392 Sang N, Fang J, Srinivas V, Leshchinsky I, Caro J. 2002. Carboxyl-terminal
2393 transactivation activity of hypoxia-inducible factor 1 alpha is governed by a
2394 von Hippel-Lindau protein-independent, hydroxylation-regulated association
2395 with p300/CBP. *Mol Cell Biol* 22(9):2984-92.
- 2396 Selak MA, Armour SM, MacKenzie ED, Boulahbel H, Watson DG, Mansfield KD,
2397 Pan Y, Simon MC, Thompson CB, Gottlieb E. 2005. Succinate links TCA
2398 cycle dysfunction to oncogenesis by inhibiting HIF-alpha prolyl hydroxylase.
2399 *Cancer Cell* 7(1):77-85.
- 2400 Semenza GL. 2007. HIF-1 mediates the Warburg effect in clear cell renal carcinoma.
2401 *J Bioenerg Biomembr* 39(3):231-4.
- 2402 Semenza GL, Jiang BH, Leung SW, Passantino R, Concordet JP, Maire P,
2403 Giallongo A. 1996. Hypoxia response elements in the aldolase A, enolase 1,
2404 and lactate dehydrogenase A gene promoters contain essential binding sites
2405 for hypoxia-inducible factor 1. *J Biol Chem* 271(51):32529-37.
- 2406 Semenza GL, Neifelt MK, Chi SM, Antonarakis SE. 1991. Hypoxia-inducible nuclear
2407 factors bind to an enhancer element located 3' to the human erythropoietin
2408 gene. *Proc Natl Acad Sci U S A* 88(13):5680-4.
- 2409 Shimomura Y, Nanaumi N, Suzuki M, Popov KM, Harris RA. 1990. Purification and
2410 partial characterization of branched-chain alpha-ketoacid dehydrogenase
2411 kinase from rat liver and rat heart. *Arch Biochem Biophys* 283(2):293-9.
- 2412 Singh VK, Hattangady DS, Giotis ES, Singh AK, Chamberlain NR, Stuart MK,
2413 Wilkinson BJ. 2008. Insertional inactivation of branched-chain alpha-keto acid
2414 dehydrogenase in *Staphylococcus aureus* leads to decreased branched-chain

- 2415 membrane fatty acid content and increased susceptibility to certain stresses.
2416 Appl Environ Microbiol 74(19):5882-90.
- 2417 Spector R, Lorenzo AV. 1973. Ascorbic acid homeostasis in the central nervous
2418 system. Am J Physiol 225(4):757-63.
- 2419 Srinivas V, Zhang LP, Zhu XH, Caro J. 1999. Characterization of an oxygen/redox-
2420 dependent degradation domain of hypoxia-inducible factor alpha (HIF-alpha)
2421 proteins. Biochem Biophys Res Commun 260(2):557-61.
- 2422 Stamford JA, Kruk ZL, Millar J. 1984. Regional differences in extracellular ascorbic
2423 acid levels in the rat brain determined by high speed cyclic voltammetry. Brain
2424 Res 299(2):289-95.
- 2425 Stolovich M, Tang H, Hornstein E, Levy G, Cohen R, Bae SS, Birnbaum MJ,
2426 Meyuhas O. 2002. Transduction of growth or mitogenic signals into
2427 translational activation of TOP mRNAs is fully reliant on the
2428 phosphatidylinositol 3-kinase-mediated pathway but requires neither S6K1
2429 nor rpS6 phosphorylation. Mol Cell Biol 22(23):8101-13.
- 2430 Strelkov AB, Fields AL, Baracos VE. 1989. Effects of systemic inhibition of
2431 prostaglandin production on protein metabolism in tumor-bearing rats. Am J
2432 Physiol 257(2 Pt 1):C261-9.
- 2433 Tang H, Hornstein E, Stolovich M, Levy G, Livingstone M, Templeton D, Avruch J,
2434 Meyuhas O. 2001. Amino acid-induced translation of TOP mRNAs is fully
2435 dependent on phosphatidylinositol 3-kinase-mediated signaling, is partially
2436 inhibited by rapamycin, and is independent of S6K1 and rpS6
2437 phosphorylation. Mol Cell Biol 21(24):8671-83.
- 2438 Terzis AJ, Niclou SP, Rajcevic U, Danzeisen C, Bjerkvig R. 2006. Cell therapies for
2439 glioblastoma. Expert Opin Biol Ther 6(8):739-49.
- 2440 Tessitore L, Costelli P, Bonetti G, Baccino FM. 1993. Cancer cachexia, malnutrition,
2441 and tissue protein turnover in experimental animals. Arch Biochem Biophys
2442 306(1):52-8.
- 2443 Visted T, Enger PO, Lund-Johansen M, Bjerkvig R. 2003. Mechanisms of tumor cell
2444 invasion and angiogenesis in the central nervous system. Front Biosci
2445 8:e289-304.
- 2446 Wang GL, Jiang BH, Rue EA, Semenza GL. 1995. Hypoxia-inducible factor 1 is a
2447 basic-helix-loop-helix-PAS heterodimer regulated by cellular O₂ tension. Proc
2448 Natl Acad Sci U S A 92(12):5510-4.
- 2449 Warburg O. 1930. The metabolism of tumors. Constable and Company, Ltd.
- 2450 Warburg O. 1956. On the origin of cancer cells. Science 123(3191):309-14.

- 2451 Whitehouse AS, Smith HJ, Drake JL, Tisdale MJ. 2001. Mechanism of attenuation of
2452 skeletal muscle protein catabolism in cancer cachexia by eicosapentaenoic
2453 acid. *Cancer Res* 61(9):3604-9.
- 2454 Wood SM, Gleadle JM, Pugh CW, Hankinson O, Ratcliffe PJ. 1996. The role of the
2455 aryl hydrocarbon receptor nuclear translocator (ARNT) in hypoxic induction of
2456 gene expression. Studies in ARNT-deficient cells. *J Biol Chem*
2457 271(25):15117-23.
- 2458 Xu G, Marshall CA, Lin TA, Kwon G, Munivenkatappa RB, Hill JR, Lawrence JC, Jr.,
2459 McDaniel ML. 1998. Insulin mediates glucose-stimulated phosphorylation of
2460 PHAS-I by pancreatic beta cells. An insulin-receptor mechanism for
2461 autoregulation of protein synthesis by translation. *J Biol Chem* 273(8):4485-
2462 91.
- 2463 Yudkoff M, Daikhin Y, Grunstein L, Nissim I, Stern J, Pleasure D. 1996. Astrocyte
2464 leucine metabolism: significance of branched-chain amino acid
2465 transamination. *J Neurochem* 66(1):378-85.
- 2466 Yudkoff M, Nissim I, Hertz L. 1990. Precursors of glutamic acid nitrogen in primary
2467 neuronal cultures: studies with ¹⁵N. *Neurochem Res* 15(12):1191-6.
- 2468 Yudkoff M, Nissim I, Kim S, Pleasure D, Hummeler K, Segal S. 1983. [¹⁵N] leucine
2469 as a source of [¹⁵N] glutamate in organotypic cerebellar explants. *Biochem*
2470 *Biophys Res Commun* 115(1):174-9.
- 2471 Zhang X, Fei Z, Bu X, Zhen H, Zhang Z, Gu J, Chen Y. 2000. Expression and
2472 significance of urokinase type plasminogen activator gene in human brain
2473 gliomas. *J Surg Oncol* 74(2):90-4.
2474
2475